

**PRECAUTIONS:** Exceeding the highest recommended daily dosage of 3000 units per pound of body weight, administering at recommended levels for more than 7 consecutive days and/or exceeding 10 mL intramuscularly per injection site may result in antibiotic residues beyond the withdrawal time.

**WARM TO ROOM TEMPERATURE AND SHAKE WELL BEFORE USING**

**WARNING:** Not for use in horses intended for food. Milk taken from animals during treatment and for 48 hours after the last treatment must not be used for food. Treatment should not exceed seven days in non-lactating dairy cattle and beef cattle, sheep and swine, or five days in lactating dairy cattle.

Discontinue use of this drug for the following time periods before treated animals are slaughtered for food: Cattle - 4 days, Sheep - 8 days, Swine - 6 days, non-ruminating calves - 7 days.

Product No.: 1PR0306  
8PR0317-908

# Pro-Pen-G<sup>®</sup> Injection

(Penicillin G Procaine Injectable Suspension)

*Aqueous Suspension  
Antibiotic  
300,000 units per mL*

*For Intramuscular Use Only  
Sterile, Multiple Dose Vial*

**Restricted Drug (California) - Use Only as Directed**

**NADA 065-505, Approved by FDA**

**Net Contents: 500 mL**

Manufactured for:  
Bimeda, Inc.  
Le Sueur, MN 56058



**Each mL of suspension contains:**

Penicillin G Potassium 300,000 units\*, Procaine hydrochloride 130.8 mg\*, Sodium citrate 10 mg, Povidone 5 mg, Lecithin 6 mg, Sodium carboxymethylcellulose 1 mg, Methylparaben 1.3 mg, Propylparaben 0.2 mg, Sodium formaldehyde sulfoxylate 0.2 mg, Procaine hydrochloride 20 mg, and Water for Injection, q.s.

\*PENICILLIN G POTASSIUM AND PROCAINE HYDROCHLORIDE REACT TO FORM PENICILLIN G PROCAINE.

FOR ANIMAL USE ONLY  
KEEP OUT OF REACH OF CHILDREN

See insert for directions and dosage.

**STORE BETWEEN 2°-8°C (36°-46°F).**  
**Avoid freezing.**

TAKE TIME



OBSERVE LABEL  
DIRECTIONS

Pro-Pen-G is a Registered Trademark of Bimeda, Inc.  
Bimeda, Inc. is a Division of Cross Vetpharm Group Ltd.

Lot No./Exp. Date:

