

NDC 57561-232-05

PROHIBIT®

(levamisole hydrochloride)
Soluble Drench Powder
Anthelmintic

Each packet contains 46.8 grams of levamisole hydrochloride activity.

FOR ORAL USE IN CATTLE AND SHEEP

Administer as a standard drench with standard drench syringe or administer as a concentrated drench solution with an automatic drenching syringe.

RECOMMENDATIONS:

Prohibit (levamisole hydrochloride) is a broad-spectrum anthelmintic and is effective against the following nematode infections in cattle and sheep:

STOMACH WORMS: (*Haemonchus*, *Trichostrongylus*, *Ostertagia*)

INTESTINAL WORMS: (*Trichostrongylus*, *Cooperia*, *Nematodirus*, *Bunostomum*, *Oesophagostomum*) [*Chabertia* - sheep only]

LUNGWORMS: (*Dictyocaulus*)

FOR ANIMAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

Restricted Drug (California) Use Only As Directed

ANADA 200-225, Approved by FDA

NET WEIGHT: 52 g (1.8 oz.)



AgriLabs®
Making A Healthy Difference®

Mfd. for: Agri Laboratories, Ltd.
St. Joseph, MO 64503

© Registered Trademark of Agri Laboratories, Ltd.

DOSAGE AND ADMINISTRATION

CATTLE--STANDARD DRENCH

SOLUTION--Place the contents of this packet in a 1 quart (32 fl. oz.) container, fill with water, swirl until dissolved. Administer as a single drench dose according to the following table:

Weight	Drench Dosage	Packet Will Treat
200 lb.	1/2 fl. oz.	64 head
400 lb.	1 fl. oz.	32 head
600 lb.	1 1/2 fl. oz.	21 head
800 lb.	2 fl. oz.	16 head

CONCENTRATED DRENCH

SOLUTION--For use with automatic syringe. Place the contents of this packet in a standard household measuring container and add water to the 8 3/4 fl. oz. level; or use the measuring container available from your supplier and add water to the mark. Swirl until dissolved. Give 2 ml (milliliter) per 100 lb. body weight. Refer to the table above for the number of cattle this packet will treat.

NOTE: Careful weight estimates are essential for proper performance of this product. Prepare solutions as needed. However, excess solutions may be stored in clean closed containers up to 90 days without loss of anthelmintic activity.

Cattle and Sheep maintained under conditions of constant helminth exposure may require re-treatment within two to four weeks after the first treatment.

RESIDUE WARNING: Do not administer to cattle within 48 hours of slaughter for food. Do not administer to sheep within 72 hours of slaughter for food. To prevent residues in milk, do not administer to dairy animals of breeding age.

CAUTION: Muzzle foam may be observed. However, this reaction will disappear within a few hours. If this condition persists, a veterinarian should be consulted. Follow recommended dosage carefully. Consult veterinarian before using in severely debilitated animals. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

Store between 15-30°C (59-86°F).

MF: L-6232-05

Rev: 5/10

LOT

EXP.

SHEEP--STANDARD DRENCH

SOLUTION--Place the contents of this packet in a 1 gallon (128 fl. oz.) container, fill with water, swirl until dissolved. Administer as a single drench dose according to the following table:

Weight	Drench Dosage	Packet Will Treat
50 lb.	1/2 fl. oz.	256 head
100 lb.	1 fl. oz.	128 head
150 lb.	1 1/2 fl. oz.	84 head
200 lb.	2 fl. oz.	64 head

CONCENTRATED DRENCH

SOLUTION--For use with automatic syringe. Place the contents of this packet in a standard household measuring container and add water to the 17 1/2 fl. oz. level. Swirl until dissolved. Give 2 ml per 50 lb. body weight. Refer to the table above for the number of sheep this packet will treat.



3