

NDC 57561-232-60

PROHIBIT®

(levamisole hydrochloride)
Soluble Drench Powder
Anthelmintic

For Oral Use in Cattle and Sheep

This bottle contains 544.5 grams of levamisole hydrochloride activity.

For Animal Use Only

Keep Out of Reach of Children

Restricted Drug (California) -

Use Only as Directed

ANADA 200-225, Approved by FDA

Net Wt: 21.34 oz (1.3 lb.) (605 g)



INDICATIONS: PROHIBIT (levamisole hydrochloride) is a broad-spectrum anthelmintic and is effective against the following nematode infections in cattle and sheep:
STOMACH WORMS: (*Haemonchus*, *Trichostrongylus*, *Ostertagia*)
INTESTINAL WORMS: (*Trichostrongylus*, *Cooperia*, *Nematodirus*, *Bunostomum*, *Oesophagostomum*)
[Chabertia-Sheep only]
LUNGWORMS: (*Dictyocaulus*)

RESIDUE WARNING: Do not administer to cattle within 48 hours of slaughter for food. Do not administer to sheep within 72 hours of slaughter for food. To prevent residues in milk, do not administer to dairy animals of breeding age.

CAUTION: Muzzle foam may be observed. However, this reaction will disappear within a few hours. If this condition persists, a veterinarian should be consulted. Follow recommended dosage carefully. Consult veterinarian before using in severely debilitated animals.

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

Store between 15°-30°C (59°-86°F).

MF# L-6232-60

Rev. 06-10

® Registered Trademark of Agri Laboratories, Ltd.

Manufactured for:
Agri Laboratories, Ltd.
St. Joseph, MO 64503

LOT NO.:

EXP. DATE:

DOSAGE AND ADMINISTRATION: When you are ready to deworm your cattle or sheep, add water to the powder in this bottle up to the 3 liter mark. Swirl to mix thoroughly before using. If any solution is left over, it may be stored for up to 3 months in this tightly capped bottle, shake well before using.

DATE WATER WAS ADDED TO THIS BOTTLE

Month Day Year

Administer as a single drench dose as follows:

CATTLE--2 mL per 100 lb. body weight

Weight	Drench Dosage	Bottle Will treat
100 lb.	2 mL	1,500 head
300 lb.	6 mL	500 head
500 lb.	10 mL	300 head
700 lb.	14 mL	214 head
1,000 lb.	20 mL	150 head

SHEEP--1 mL per 50 lb. body weight

Weight	Drench Dosage	Bottle Will Treat
50 lb.	1 mL	3,000 head
100 lb.	2 mL	1,500 head
150 lb.	3 mL	1,000 head
200 lb.	4 mL	750 head

NOTE: Careful weight estimates are essential for proper performance of this product.

Cattle and sheep maintained under conditions of constant helminth exposure may require retreatment within two to four weeks after the first treatment.

