



Zinc Phosphide Rodent Pellets AG

**Kills Moles* and Gophers
Kills Rats and Mice**

This product may only be used to control Norway rats, roof rats and house mice inside and within 100 feet of buildings or inside of transport vehicles (ships, trains, or aircraft).

For Control of Pocket Gophers and Moles* in Below-Ground Applications Only

Do Not Sell This Product In Individual Containers Holding Less Than Four Pounds of Bait.

ACTIVE INGREDIENT:

Zinc Phosphide.....2.00%

OTHER INGREDIENTS:.....98.00%

TOTAL:.....100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

EPA Reg. No. 61282-56

EPA Est. No. 61282-WI-01

FIRST AID

HAVE PRODUCT LABEL WHEN OBTAINING TREATMENT ADVICE

Any person applying zinc phosphide products and experiencing signs and symptoms such as nausea, abdominal pain, tightness in the chest, or weakness, needs to be seen by a physician immediately.

If Swallowed:	<ul style="list-style-type: none"> Call a poison control center, doctor, or 1-800-498-5743 immediately for treatment advice or transport the patient to the nearest hospital. Do not give any liquid to the patient. Do not administer anything by mouth. Do not induce vomiting unless told to do so by the poison control center or doctor.
If on Skin or on Clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.

FIRST AID (cont.)

If Inhaled:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing call 911 or an ambulance, then give artificial respiration preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
If in Eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice.
Have product container or label with you when calling a poison control center or doctor for treatment advice.	
TREATMENT FOR PET POISONING: If animal eats bait, call veterinarian at once.	
NOTE TO PHYSICIAN OR VETERINARIAN: Contains the Phosphine-producing active Zinc Phosphide. Probable mucosal damage may contraindicate the use of gastric lavage. For animals ingesting bait and/or showing poisoning signs, induce vomiting by using hydrogen peroxide. Sodium bicarbonate can be given orally to neutralize the stomach acidity. The stomach and intestinal tract can be evacuated, oxygen administered and cardiac and circulatory stimulants given.	
For 24-hour emergency information on this product, call 1-800-498-5743 (US & Canada) or 1-651-523-0318 (all other areas).	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, absorbed through skin, or inhaled. Avoid breathing dust. Avoid contact with skin or clothing. Wear pants, shoes, socks, and barrier laminate gloves. Wash hands thoroughly with soap and water before eating, drinking, and chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse. Any person who retrieves carcasses or unused bait following application of this product must wear barrier laminate gloves.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing them. As soon as possible, wash hands thoroughly after applying the bait and before eating, drinking, chewing gum, using tobacco or using the toilet and change into clean clothing

ENVIRONMENTAL HAZARDS

This product is extremely toxic to birds, fish and other wildlife. Wildlife feeding on treated bait may be killed. Dogs, cats and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions, use restrictions and precautions.

Uses in Lawns to Control Moles and Pocket Gophers

not approved for the control of Moles in the following states:

California, Hawaii, Indiana, and New Hampshire

For use only against the eastern mole (*Scalopus aquaticus*) and the hairy-tailed mole (*Parascalops breweri*) in North Carolina. Not for use against the star-nosed mole (*Condylura cristata*) in North Carolina. Not for use against gophers in North Carolina.

USE RESTRICTIONS:

- This product may only be applied at the use sites and by the application methods indicated on this label.
- Bait must be applied directly into mole and pocket gopher burrow systems.
- Any person who retrieves carcasses or unused bait following application of this product must wear barrier laminate gloves.

- This product may only be used to control moles and pocket gophers in manual below ground applications.
- Do not apply in gardens where food or feed may become contaminated.
- Do not apply this product by use of an artificial burrow builder.
- Do not use this product for sod production.

SPECIFIC DIRECTIONS FOR MOLES: The presence of moles may be indicated by a network of surface ridges in the turf, by a series of conical mounds of earth pushed up from deep burrows, or by damage to flower bulbs.

Surface Ridges: Prior to treatment, determine which burrows are active according to one of the following two methods. Either press down a small section of the tunnel or remove a one inch section of the tunnel's roof. Mark these sections and recheck in 24 hours. The burrow is considered active if the flattened runway has been raised up or if the roof has been repaired. Only treat active burrows. Carefully, with a long-handled spoon, punch a hole in the top of each active tunnel, insert a teaspoon amount of this product, and carefully close the hole with sod or stone. Do not collapse the runway again or allow loose soil to cover the bait. After a few days, check to see if the burrows are still active according to one of the two methods described above. If so, repeat treatment.

Conical Mounds: With a long handled spoon, carefully remove the center plug from each conical mound. Drop a teaspoon amount of this product into the underground tunnel. Close the opening with a rock or sod. Do not allow loose dirt to cover bait. If mole activity persists after one treatment, a second treatment may be made. If mole activity persists after two treatments, abandon use of this product and seek another method for controlling moles.

Flower Bulb Protection (Fall-planted): Pack about one-quarter inch of soil against each flower bulb. Place several pellets at sides of and around the flower bulb. Then cover the bulb with soil.

Follow-Up: Wearing barrier laminate gloves, collect and dispose of all dead, exposed animals and dispose of leftover bait in accordance with disposal instructions. Use detergent and hot water to wash spoon used for baiting. Do not use this spoon for mixing, holding or transferring food or feed. If reinfestation does occur, repeat treatment

SPECIFIC DIRECTIONS FOR POCKET GOPHERS: Burrowing pocket gophers (*Thomomys* spp. and *Geomys* spp.) throw out low, fan-shaped mounds on either side of their underground tunnels. The lateral tunnels coming to the surface are on the flat sides of the fans and are plugged with soil. With a long-handled spoon, carefully remove the plug. Insert a teaspoon of this product as far down the lateral tunnel as possible. Reclose the opening, being careful not to cover bait with soil.

FOLLOW-UP: Wearing barrier laminate gloves, collect and dispose of all dead, exposed animals and dispose of leftover bait in accordance with disposal instructions. Use detergent and hot water to wash spoon used for baiting. Do not use this spoon for mixing, holding or transferring food or feed. If reinfestation does occur, repeat treatment.

Uses Associated With Buildings and Structures To Control Rats and Mice

IMPORTANT: Do not expose children, pets or other non-target animals to rodenticides.

To help to prevent accidents:

- Store product not in use in a location out of reach of children and pets.
- Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
- Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used wherever children, pets, non-target mammals, or birds may have access to the bait placement location.

FOR NORWAY RATS, ROOF RATS AND HOUSE MICE USE RESTRICTIONS

This product can only be used to control Norway rats, roof rats, and house mice in

**NET CONTENTS: 12 pounds
NET WEIGHT: 12 pounds**

and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port or terminal buildings and related structures around and associated with these sites. Fence and perimeter baiting, beyond 100 feet from any structure as defined above, is prohibited. This product must not be applied directly to food or feed crops. Do not use this product in sewers.

Do not broadcast bait. Do not place bait near or inside ventilation duct openings. Do not contaminate water, food, feed, food- or feed-handling equipment, or surfaces that come into direct contact with food. When used in USDA inspected facilities, this product must be applied in tamper-resistant bait stations.

Do not sell this product in individual containers holding less than four (4) pounds of bait.

Restrictions for Baiting Rat Burrows

This product may only be applied to active rodent burrows to control Norway rats and roof rats within or beyond 100 feet of buildings and man-made structures, provided that infestation by Norway rats or roof rats has been confirmed. Bait must be placed no less than 6 inches into active Norway/roof rat burrows.

Because Norway/roof rat infestations may occur in areas farther than 100 feet from buildings and man-made structures when the rodents have ample supplies of food and cover, efforts should be made to remove conducive food trash, garbage, clutter, and debris.

SELECTION OF TREATMENT AREAS: After removing as much alternate food as possible, determine dry, acid-free areas where rats or house mice will most likely find and consume the bait. Generally these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where signs of rats or house mice have been observed.

APPLICATION DIRECTIONS

Rats: Apply 1 to 2 teaspoon amounts of bait per placement. For outdoor use, place bait in active rat burrows or tamper-resistant bait stations. Maintain an uninterrupted supply of fresh bait until there no longer are signs of new feeding by rats. Do not treat the same area at less than 30 day intervals.

Mice: Apply 1 teaspoon amount of bait at each placement location. Space placements at 8- to 12-foot intervals. For outdoor use, place bait in tamper-resistant bait stations. Maintain an uninterrupted supply of fresh bait until there are no longer signs of new feeding by house mice.

Follow-Up: Wearing barrier laminate gloves, collect and dispose of all dead, exposed animals and dispose of leftover bait in accordance with disposal instructions. Use detergent and hot water to wash spoon used for baiting. Do not use this spoon for mixing, holding or transferring food or feed. If reinfestation does occur, repeat treatment.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE:

Store in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

PESTICIDE DISPOSAL:

Pesticide wastes are toxic. Improper disposal of excess pesticide is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

Non-refillable container. Do not reuse or refill this container. Dispose of empty container by placing in trash, or at an approved waste disposal facility

DISCLAIMER

DISCLAIMER: To the extent permitted by applicable law, seller makes no representation or warranty, either express or implied, for results due to misuse, improper handling or improper storage of this material. Nor does Seller assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or improper storage of this material.

For non-emergency (e.g., current product information) call: 1-800-621-8829


*Not Approved of this Use Pattern in the Following States: California, Hawaii, New Hampshire, and Indiana.



Item No. 114600

Manufactured By:
HACCO, Inc.
110 Hopkins Dr.
Randolph, WI 53956

Code (NO.RA.) NO.RA
LA 220069



0 23626 01195 3