

**INDICATIONS**

**BROILER AND REPLACEMENT CHICKENS:** Use for the treatment of disease outbreaks of coccidiosis, fowl cholera, and infectious coryza.

**MEAT PRODUCING TURKEYS:** Use for the treatment of disease outbreaks of coccidiosis and fowl cholera.

**DAIRY CALVES, DAIRY HEIFERS AND BEEF CATTLE:** Use for the treatment of shipping fever complex and bacterial pneumonia associated with *Pasteurella* spp. sensitive to sulfadimethoxine, calf diphtheria and foot rot associated with *Fusobacterium necrophorum* (*Sphaerophorus necrophorus*) sensitive to sulfadimethoxine.

**PRECAUTIONS**

Store at controlled room temperature 20° to 25°C (68° to 77°F), excursions permitted from 15° to 30°C (59° to 86°F). If freezing occurs, thaw before using. Protect from light; direct sunlight may cause discoloration. Freezing or discoloration does not affect potency. Prepare a fresh stock solution daily.

**CHICKENS AND TURKEYS:** If animals show no improvement within 5 days, discontinue treatment and re-evaluate diagnosis. Handle the recommended dilutions (chickens 0.05% and turkeys 0.025%) as regular drinking water. Administer as sole source of drinking water and sulfonamide medication.

Chickens and turkeys that have survived fowl cholera outbreaks should not be kept for replacements or breeders.

**CATTLE:** During treatment period, make certain that animals maintain adequate water intake. If animals show no improvement within 2 or 3 days, re-evaluate diagnosis. Treatment should not be continued beyond 5 days.



FOR USE IN ANIMALS ONLY  
NOT FOR HUMAN USE  
KEEP OUT OF REACH OF CHILDREN  
RESTRICTED DRUG (California) - USE ONLY AS DIRECTED

**RESIDUE WARNINGS**

**CHICKENS AND TURKEYS:** Withdraw 5 days before slaughter. Do not administer to chickens over 16 weeks (112 days) of age or to turkeys over 24 weeks (168 days) of age.

**CATTLE:** Withdraw 7 days before slaughter.

For dairy calves, dairy heifers and beef cattle only. A withdrawal period has not been established for this product in pre-ruminating calves.



DO NOT USE IN CALVES  
TO BE PROCESSED FOR VEAL

Strategic Veterinary Pharmaceuticals, Inc.  
1411 Oak Street  
Elwood, KS 66924

LBL70001-003, R3  
Revision Date 07-16

Lot No. \_\_\_\_\_

Exp. Date: \_\_\_\_\_



## SDM Sulfadimethoxine Concentrated Solution 12.5%

**CAUTION:** Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

**ANTIBACTERIAL**  
for Use in Drinking Water

For Oral Use in Chickens,  
Turkeys, and Cattle

ANADA 200-165, Approved by FDA

**NET CONTENTS: 1 GALLON (3.785 liters)**

**PROTECT FROM LIGHT**

3.75 g sulfadimethoxine/fl oz.

**Dosage and Administration****Chickens and Turkeys**

**Treatment Period:** 6 consecutive days

**Recommended Concentrations:** Chickens – 0.05%, Turkeys – 0.025%

**Chickens** Add 1 fl oz (30 mL) to 2 gallons of drinking water -or- 25 fl oz to 50 gallons of drinking water.

**Turkeys** Add 1 fl oz (30 mL) to 4 gallons of drinking water -or- 25 fl oz to 100 gallons of drinking water.

**Automatic Proportioners\* Stock Solution –** To make 2 gallons of Stock Solution use:

**Chickens** Add 1 gallon of SDM Concentrated Solution 12.5% to 1 gallon of water

**Turkeys** Add 2 qts of SDM Concentrated Solution 12.5% to 6 qts of water

\*Set proportioner to a feed rate of 1 fl oz (30 mL) of SDM (Sulfadimethoxine) Stock Solution per gallon of water.

**Dairy Calves, Dairy Heifers, and Beef Cattle**

**Treatment Period:** 5 consecutive days

**Dosage:** Initial dose of 25 mg/lb followed by four maintenance doses of 12.5 mg/lb.

**Summer Administration**

Dosage recommendations for summer are based on an estimated water intake of 1 gallon of water per every 100 lb of body weight per day.

**SDM Concentrated Solution 12.5%**

Daily Drinking Water Supply	Initial Dose	Maintenance Dose
25 gal	1 pt (16 fl oz)	1 cup (8 fl oz)
50 gal	1 qt (32 fl oz)	1 pt (16 fl oz)
200 gal	1 gal (128 fl oz)	2 qt (64 fl oz)

**Winter Administration**

Dosage recommendations for winter are based on an estimated water intake of 1 gallon of water per every 150 lb of body weight per day.

**SDM Concentrated Solution 12.5%**

Daily Drinking Water Supply	Initial Dose	Maintenance Dose
16 gal	1 pt (16 fl oz)	1 cup (8 fl oz)
33 gal	1 qt (32 fl oz)	1 pt (16 fl oz)
127 gal	1 gal (128 fl oz)	2 qt (64 fl oz)

**For individual treatment of cattle,** SDM Concentrated Solution 12.5% may be given as a drench. Administer using an initial dose of 25 mg/lb followed by 4 maintenance doses of 12.5 mg/lb/day. One fl oz will medicate one 150-lb animal initially and 1/2 fl oz will medicate one 150-lb animal on maintenance dose.