

# Sacox<sup>®</sup>60

## Brand of salinomycin sodium Type A Medicated Article

For use in broiler, roaster, replacement  
(breeder and layer) chickens and quail only.

**Active drug Ingredient** Salinomycin equivalent to 60 grams salinomycin sodium activity per pound.

**For use in manufacturing medicated feeds only**

**Broiler, roaster and replacement (breeder and layer) chickens** Use Sacox<sup>®</sup> 60 to provide salinomycin sodium concentrations ranging from 40-60 g/ton (0.0044% to 0.0066%) for the prevention of coccidiosis in broiler, roaster and replacement chickens caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti* and *E. mixti*. The dosage should be adjusted to meet the severity of the coccidial challenge, which varies with environmental and management conditions.

**Quail** Use Sacox<sup>®</sup> 60 to provide salinomycin sodium concentrations of 50 g/ton (0.0055%) for the prevention of coccidiosis in quail caused by *Eimeria dispersa* and *E. lettyae*.

**Mixing Directions** Thoroughly mix Sacox<sup>®</sup> 60 with non-medicated feed to provide the level of salinomycin sodium per ton as indicated in the table below.

Salinomycin Sodium Activity (grams per ton)	Amount of Sacox <sup>®</sup> 60 Type A Medicated Article (60 g/lb) Per Ton of Feed	
	Lbs	Grams
40	0.67	304
45	0.75	341
50	0.83	377
55	0.92	418
60	1.00	454

**Feeding Directions** Feed continuously as the only ration.

**Important** Must be thoroughly mixed in feeds before use. Not approved for use with pellet binders.

**Caution** May be fatal if accidentally fed to adult turkeys or to horses.

**Warning: Do not feed to laying hens producing eggs for human consumption. When mixing and handling Sacox<sup>®</sup> 60 premix, use protective clothing, impervious gloves, eye protection and an approved dust respirator. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water.**

STORAGE: Store at or below 25°C (77°F), excursions permitted to 40°C (104°F). Avoid prolonged exposure to high humidity.

ANADA #200-075, Approved by FDA

Distributed by:  
**HUVEPHARMA, INC.**  
Peachtree City, GA 30269

5042-06 Rev. 05-14



Net wt. 50 lb (22.68 kg)

# Sacox<sup>®</sup>60

## Brand of salinomycin sodium Type A Medicated Article

**Active Drug Ingredient:**  
Salinomycin equivalent to 60 grams  
salinomycin sodium activity per pound.

**Other Ingredients:**  
Calcium Carbonate, Mineral Oil,  
Calcium Silicate and Silicon Dioxide.

Important: See back panel for use  
directions.

Not For Human Use  
Restricted Drug (California) - Use only as Directed

Net Weight 50 lb (22.68 kg)



(salinomycin sodium)  
**Sacox<sup>®</sup>60**

229528 H-236-003 7-12  
FREIGHT SHIPPING BAG  
MEETS REQUIREMENTS OF  
APPLICABLE FREIGHT CLASSIFICATION  
QUANTITY:  
BANCROFT BAG, INC.  
WEST HAVEN, CT 06457-1000  
16x5.5x32

**Sacox<sup>®</sup>60**  
(salinomycin sodium)

Pre-Distorted Flexo Template  
PBOM 16" x 5.5" x 32" Print Length-32"  
Factory Close Bottom to Back .067  
BBI-BMC 4-16-08

