

SANTOQUIN[®] LIQUID

Feed and Forage Antioxidant

GUARANTEED ANALYSIS

Ethoxyquin (Minimum)..... 91.0%
p-Phenetidine <3.0%
Specific Gravity1.030 @25°C

INGREDIENT

Ethoxyquin

THIS PRODUCT IS FOR LIVESTOCK FEED AND FEED INGREDIENTS ONLY.

Product shelf life in sealed containers is 3 years from date of manufacture.

Store product in original container at room temperature.

Do not store below 0°C or above 50°C.

DIRECTIONS FOR USE

The viscosity of this product will vary inversely with temperature.

FISHMEAL: As a preservative for fishmeal that is intended for use in feeds, use SANTOQUIN[®] Liquid at a level of 400 to 1000 parts per million, depending on the oil content of the fish. The fishmeal label must show the ethoxyquin concentration, and directions for use to ensure that the finished feed contains no more than 0.015% ethoxyquin.

LIVESTOCK FEED: Ethoxyquin, a chemical preservative, is used for retarding oxidative destruction of carotene, xanthophylls and vitamins A and E in feeds.

Recommended uses include addition as a preservative mixed in the final feed, inclusion as an ingredient in premixes and concentrates, and addition to suitable fats or other liquids prior to addition to the feed.

Add SANTOQUIN[®] Liquid at a rate of 0.0125% or 0.125 kg per metric ton of final manufactured feed. The total amount of the SANTOQUIN[®] Liquid added plus any furnished from other feed ingredients treated with ethoxyquin must not exceed 0.015% of the final mixed feed. Intermediate premixes and concentrates should clearly indicate the concentration level and include directions for use of the pre-mix or concentrate to insure that the final mixed feed will contain no more than the approved level of use.


DEHYDRATED FORAGE CROPS: Ethoxyquin is used as a chemical preservative for the purpose of retarding oxidative destruction of naturally occurring carotenes and vitamin E in forage crops. Ethoxyquin is added to the dehydrated forage crops in an oil mixture containing only feed grade fat or oil after dehydration and/or prior to grinding and mixing. Add SANTOQUIN[®] Liquid at a rate of 0.0150% or 0.15 kg per metric ton of dehydrated forage.

TOTAL ETHOXYQUIN LEVELS: In finished feeds, total ethoxyquin from SANTOQUIN[®] Liquid and any other source of ethoxyquin must not exceed 0.015% (150 parts per million).

Distributed and guaranteed by: EW Nutrition USA, Inc., 28800 R Avenue, Adel, Iowa 50003 United States of America Phone: +01 515 478 1901

 EW Nutrition GmbH
Hogenbögen 1, 49429 Visbek, Germany
α DE NI 400130
Phone: + 49 4445 9868 0
www.ew-nutrition.com

**Net Weight: 1000 kg
(2204.6 lb)**



WARNING

Harmful if swallowed. May cause an allergic skin reaction. Suspected of causing genetic defects.

CAUTION

Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Avoid breathing mist or vapor. Wash thoroughly after handling. Do not eat, drink or smoke when using this product. Contaminated work clothing must not be allowed out of the workplace. Wear protective gloves/ protective clothing/eye protection /face protection

If exposed or concerned: Get medical advice/attention. If swallowed: Call a poison center/doctor if you feel unwell. Rinse mouth. If on skin: Wash with plenty of water. If skin irritation or rash occurs: Get medical advice/attention. Take off contaminated clothing and wash before reuse.

LOT NUMBER:

MANUFACTURE DATE:

EXPIRATION DATE:

80001417v9