

INDICATIONS:

For use in pigs under four (4) weeks of age for treatment and control of porcine enteric colibacillosis (scours) caused by *E. coli* susceptible to spectinomycin.

EACH mL CONTAINS:

Spectinomycin (from spectinomycin dihydrochloride pentahydrate) 50 mg

WARNING:

This product is only intended for use in pigs under four (4) weeks of age or weighing less than 15 lbs.

Do not administer within 21 days of slaughter.

Store at controlled room temperature, 20°-25° C (68°-77°F)



SpectoGard and Scour-Chek are Registered Trademarks of Bimeda, Inc.

Manufactured by: Bimeda-MTC Animal Health Inc.
Cambridge, Ontario, Canada N3C 2W4

Bimeda, Inc. and Bimeda-MTC Animal Health Inc.
are Divisions of Cross Vetpharm Group Ltd.

SpectoGard®

Scour-Chek®

(Spectinomycin)

ORAL SOLUTION FOR PIG SCOURS

ANTI-INFECTIVE

FOR ANIMAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

Restricted Drug - Use Only As Directed (California)

ANADA 200-364, Approved by FDA

NET CONTENTS:

1000 mL (33.8 fl oz)

Manufactured for:
Bimeda, Inc.
Le Sueur, MN 56058



Bimeda

DOSE:

Pigs under 10 lbs. - 1 pump (1 mL) twice daily.
Pigs over 10 lbs. - 2 pumps (2 mL) twice daily.
Each pump of the plunger delivers 1 mL of solution containing 50 mg of spectinomycin. Treatment may be continued twice daily for 3 to 5 days. If pigs do not improve within 48 hours, rediagnosis is suggested.

DIRECTIONS FOR USE:

The 1000 mL bottle of SpectoGard® Scour-Chek® should be used only to refill the 240 mL container in the following manner.

1. Remove the cap and doser from the 240 mL bottle.
2. Unscrew the cap from the 1000 mL bottle.
3. Pour, from the 1000 mL bottle, the volume desired.
4. Reinsert the doser in the 240 mL bottle and screw the top on tightly.
5. Screw the top tightly on the 1000 mL bottle.

Follow the detailed instructions for use shown on the carton containing the 240 mL bottle of SpectoGard® Scour-Chek®.

Product No. 1SPE011
8BTL104

UPC 80%
8 99855 60049 3