



011 1 03 44104 99100 0

# Stafac<sup>®</sup> 50 (virginiamycin)

TYPE A  
MEDICATED ARTICLE

(50 g virginiamycin activity per lb)  
For use in swine and poultry feeds as specified below



#### Directions for Use Mixing Directions—Swine

Stafac 50 Type A Medicated Article must be thoroughly mixed in the feed to assure even distribution. Determine the appropriate amount of Stafac 50 required from the table below. Dilute Stafac 50 with a portion of one of the feed ingredients (enough to make a quantity equal to 10 lb per ton) before final mixing.

	Required level of virginiamycin activity per ton of Type C Medicated Feed (complete feed)	Required amount of Stafac 50 per ton of Type C Medicated Feed (complete feed)	Amount of virginiamycin activity provided to animal per lb of body weight per day
Swine			
Treatment of swine dysentery in nonbreeding swine weighing over 120 lb	100 g for 2 weeks	2 lb	2.27 mg
Treatment and control of swine dysentery in swine weighing up to 120 lb	100 g for 2 weeks followed by 50 g	2 lb 1 lb	2.27 mg 1.14 mg
Aid in control of swine dysentery in swine weighing up to 120 lb	25 g	8 oz	0.57 mg

For use in animals or on premises with a history of swine dysentery but where symptoms have not yet occurred.

CAUTION: Not for use in breeding swine over 120 lbs.

#### Mixing Directions—Poultry

Mix 1 lb of Stafac 50 with 9 lb of feed ingredients to make 10 lb of an intermediate mix to contain 5 g/lb virginiamycin. Add 1 lb of the intermediate feed premix for each 5 g of virginiamycin needed to medicate a ton of feed.

	Required level of virginiamycin activity per ton of Type C Medicated Feed (complete feed)	Required amount of Stafac 50 per ton of Type C Medicated Feed (complete feed)
Poultry		
For use in broiler chickens		
Prevention of necrotic enteritis caused by <i>Campylobacter parvipes</i> susceptible to virginiamycin	20 g	180 g

Not for Use in Laying Chickens

Feed Continuously as Sole Ration  
Store At Or Below 25°C (77°F); Excursions Permitted Up To 40°C (104°F)  
Close Container After Use  
Not For Human Use  
Restricted Drug (California) — Use Only as Directed

(virginiamycin)  
**Stafac<sup>®</sup> 50**

Phibro  
ANIMAL HEALTH CORPORATION



# Stafac<sup>®</sup> 50 (virginiamycin)

TYPE A  
MEDICATED ARTICLE

#### Active Drug Ingredient

Virginiamycin ..... 11%  
(Contains 50 g virginiamycin activity per lb)

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

#### Ingredients:

Processed grain by-products, roughage products, calcium carbonate, carboxymethylcellulose, mineral oil

For further manufacture only.

Follow Directions on Back Panel

Stafac is a registered trademark of Phibro Animal Health Corporation  
Phibro Animal Health Corporation, Teaneck, NJ 07666

NADA #91-467, Approved by FDA  
9910000  
101-9107-09A

Net Weight 50 lb (22.6 kg)

**Phibro**  
ANIMAL HEALTH CORPORATION

Phibro  
ANIMAL HEALTH CORPORATION

**Stafac<sup>®</sup> 50**  
(virginiamycin)

9910000  
101-9107-09A

Phibro  
Phibro Animal Health Corporation  
Teaneck, NJ 07666