



Cyanarox[®] Insecticidal Bait

Granular insect bait for use in and around residential, commercial, institutional and agricultural structures to control listed fly pests and kill listed cockroaches

- ★ Kills Houseflies, Stable flies, and other nuisance flies
- ★ Kills Oriental, German, and American cockroaches

SPECIMEN LABEL

ACTIVE INGREDIENT:

Cyantraniliprole*
 3-bromo-1-(3-chloro-2-pyridinyl)-N-[4-cyano-2-methyl-6-[(methylamino)carbonyl]phenyl]-1H-pyrazole-5-carboxamide:0.5%

OTHER INGREDIENTS:.....99.5%

TOTAL:.....100.0%

*Cyantraniliprole belongs to the anthranilic diamide chemical class
 Ready-to-use granular insect bait containing 0.005 lb cyantraniliprole per pound of product.

KEEP OUT OF REACH OF CHILDREN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
 (If you do not understand the label, find someone to explain it to you in detail.)

CYANTRANILIPROLE	GROUP	28	INSECTICIDE
------------------	-------	----	-------------

EPA Reg.No. 100-1541-89459
 EPA Est. No. 39578-TX-1

Starbar[®] Cyanarox[®] Insecticidal Bait (Cyanarox[®] Insecticidal Bait) is a ready-to-use granular insect bait containing 0.005 lb cyantraniliprole per pound of product.

FIRST AID

HOT LINE NUMBER

For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire or Accident), Call 1-800-347-8272.

PRECAUTIONARY STATEMENTS

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear: • Shirt and long pants, • Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Runoff may be hazardous to aquatic organisms in water adjacent to use sites.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read and understand the entire label before using this product.

Cyanarox[®] Insecticidal Bait must be used in accordance with the Directions For Use on this label, or exemptions under FIFRA (Special Local Need Registrations, FIFRA Section 18 exemptions, FIFRA 2(ee) Bulletins). Always read the entire label, including the Limitation of Warranty and Liability.

PRODUCT INFORMATION: Cyanarox[®] Insecticidal Bait contains the active ingredient cyantraniliprole in a ready-to-use granular bait formulation:

- Controls listed fly pests and kills cockroaches in and around residential, commercial, institutional and agricultural structures;
- Provides an alternative mode of action for insecticide resistance management;
- Can be an important tool in an integrated pest management program.

APPLICATION USE SITES: Cyanarox[®] Insecticidal Bait may be applied in these areas:

- outdoor waste-storage areas around commercial establishments, such as restaurants, taverns, hotels, grocery stores, supermarkets, bakeries, warehouses, and sports complexes;

- outdoor areas of commercial operations, such as canneries; beverage-processing plants; dairy-, meat-, and poultry-processing plants; fruit- and vegetable-processing plants; seafood-processing plants; tanneries; rendering plants; grease-collection stations; kennels; schools; recycling plants; and military installations;
- outdoor areas of agricultural animal-production facilities, such as poultry/broiler houses, swine-production structures, livestock-housing structures, and horse stables.

To control house fly, stable fly, dump fly, fruit flies, dung fly, and black-eyed fly, and to kill cockroaches, Cyanarox® Insecticidal Bait may be used indoors at the sites listed above as a water-diluted application. For flies only, Cyanarox® Insecticidal Bait may be applied as a granular if placed in bait stations. In walkways within caged-layer houses or in penned-animal facilities, Cyanarox® Insecticidal Bait may be applied as a broadcast, scatter bait. Refer to the TARGET PESTS table describing **Use in Bait Stations** or **Alternative Application Methods** for specific use instructions.

USE RESTRICTIONS AND PRECAUTIONS

- **DO NOT** apply as a scatter bait indoors in residential structures, commercial establishments, food- and feed-handling establishments, food-processing plants, restaurants and animal-production facilities. Bait stations or the alternative application methods (see following section) must be used when applying Cyanarox® Insecticidal Bait inside these buildings/structures, except for walkways within caged-layer houses or penned-animal facilities.
- **DO NOT** apply as a scatter bait in areas accessible to children and pets. When bait stations or the alternative application methods (see following section) are used, place stations, ropes, or cards out of the reach of children and pets.
- **DO NOT** place bait within reach of food-producing animals or animals being grown for food.
- **DO NOT** apply as a scatter bait outdoors when raining.
- **DO NOT** use outdoors as a scatter bait in the Nassau, Suffolk, Kings, and Queens Counties of New York State.
- For outdoor broadcast applications, **DO NOT** apply more than 2 lb per 1,000 sq ft per calendar year (equivalent to 0.01 lb ai per 1,000 sq ft per calendar year or 0.44 lb ai/A per calendar year).

Resistance Management:

CYANTRANILIPROLE	GROUP	28	INSECTICIDE
------------------	-------	----	-------------

Cyanarox® Insecticidal Bait is a Group 28 Insecticide that contains the active ingredient cyantraniliprole. Due to the inherent risk of the development of resistance to any pesticide, it is strongly advised that Cyanarox® Insecticidal Bait be used in a sound resistance-management program that includes, but is not limited to:

- Applying Cyanarox® Insecticidal Bait according to the label-use directions;
- Alternating Cyanarox® Insecticidal Bait with products with different modes of action;
- Avoiding treatment of successive generations with Cyanarox® Insecticidal Bait.

For additional information about implementing these or other resistance management practices, consult a local or state Extension specialist or Central Garden & Pet representative.

Food- and Feed-Handling Establishments: (Food- and feed-handling establishments are places other than private residences in which food and feed are held, processed, prepared or served.)

- For indoor areas of residential, institutional, public, agricultural buildings/structures, or food- and feed-handling establishments, Cyanarox® Insecticidal Bait must be placed in bait stations suitable for granular insect baits or prepared and applied as a water-diluted product. Indoor use as a scatter bait is permitted only on walkways in caged-layer houses or penned-animal facilities.
- **DO NOT** apply to areas or surfaces where food, feed, food utensils or food-processing surfaces may come into contact and become contaminated.

Applications in Non-food and Non-feed Areas: Cyanarox® Insecticidal Bait is approved for use to control house flies and other listed fly pests and to kill cockroaches in residential structures and the non-food and non-feed areas of commercial, industrial, public, agricultural and institutional buildings/structures, including restaurants, warehouses, food- and feed-processing plants, supermarkets, hospitals, nursing homes, motels, hotels, schools, daycares, laboratories, computer facilities, aircraft, buses, boats/ships, trains, pet shops and zoos.

Non-food and non-feed areas of food- and feed-handling establishments include areas such as garbage rooms, lavatories, floor drains (to sewers), entries, and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after bottling or canning).

Refer to section below for use directions and restrictions when making applications in food/feed handling areas of Food- and Feed-Handling Establishments.

Applications in Food- and Feed-Handling Areas: Application to food/feed areas of food- and feed-handling establishments may only be made using bait stations suitable for use with granular insect control baits.

Food- and feed-handling areas include areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing edible waste storage, and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas are also considered a food and feed area when food is exposed and facility is in operation.

APPLICATION PROCEDURES AND USE RATES

Broadcast or scatter Cyanarox® Insecticidal Bait using suitable granular spreader equipment or directly from the container evenly to the target areas. Avoid placing the bait granules in piles and scatter bait as evenly as possible over surfaces.

Apply 0.2 lb (3.2 oz) of Cyanarox® Insecticidal Bait per 1,000 sq ft when listed fly and cockroach pest populations are low. Apply 0.4 lb (6.4 oz) of bait per 1,000 sq ft when populations are high. Reapply bait every 7 days until insect pest population is at acceptable levels. For outdoor broadcast applications, **DO NOT** apply more than 2 lb per 1,000 sq ft per calendar year.

Use in Bait Stations: Use bait stations that are suitable for granular baits. Apply 0.2 lb (3.2 oz) of Cyanarox® Insecticidal Bait when listed fly pest populations are low using enough bait stations to make the necessary amount of bait available per 1,000 sq ft. Apply 0.4 lb (6.4 oz) of bait in bait stations per 1,000 sq ft when listed fly pest populations are high. Replenish bait stations until the fly population is at acceptable levels in the target area.

Alternative Application Methods (for Listed Pests)

Application of Cyanarox® Insecticidal Bait to Glue-Boards or Sticky Cards:

Cyanarox® Insecticidal Bait may be applied as a dry scatter bait to the sticky surfaces of glue boards or other sticky cards as a way to present granular bait to listed fly and cockroach pests. Apply Cyanarox® Insecticidal Bait to sticky surface of cards in a way so that the sticky material is covered with an even layer of bait (approximately 0.1 oz per sq inch of card surface). The amount used per card depends on the size and type of sticky surface.

Hang cards from ceiling supports or other means to provide a treated surface for flies to land. Replace cards as bait is consumed or if cards become dirty, but not more frequently than every 7 days.

Use of Water-Diluted Cyanarox® Insecticidal Bait Product – How to Prepare Dilution:

Cyanarox® Insecticidal Bait may be applied by diluting granules in warm water at ratios ranging from 0.5-1.5 oz Cyanarox® Insecticidal Bait per 1 fl oz water. Mix in non-food use container and stir until bait has dissolved. Let mixture stand until a suitable consistency is achieved that allows one to paint the mixture onto a surface or apply to disposable cards or rope wicks.

Methods of Applications of Water-Diluted Product (not for Stable Flies)

- **Paint-on method.** Prepare water-diluted product as described above. Apply as a spot treatment or in 6-12 inch wide bands to surfaces every 10 ft where flies are likely to rest using a paint brush, roller or other applicator suited to spreading water-diluted bait onto a surface. Rinse applicator thoroughly after using. Reapply diluted bait as needed, but not more frequently than every 7 days. Some surfaces may become temporarily discolored when painting with diluted bait so it is advisable to test paint a small area to determine surface compatibility.
- **Application to disposable cards.** Water-diluted bait may also be applied to cards. Hang cards from ceiling supports or other means to provide a treated surface for flies to land. Treated cards may be disposed of when they are no longer attractive to pests or when they become dirty and may be replaced with freshly treated cards every 7 days.
- **Rope-wick application of water-diluted product.** Rope wick may be used as a disposable surface for exposing flies to water-diluted Cyanarox® Insecticidal Bait.

Prepare water-diluted product as described above in a container. Place short pieces of rope-wick (approximately 6-36 inch lengths) into diluted

product and allow to soak up diluted bait for 3-5 minutes. Remove rope pieces, allow to dry, and hang in areas where flies are likely to rest.

Surfaces that may be treated with paint-on treatments or where disposable boards/cards and/or rope wicks may be hung include doors and windows, building perimeter foundations, walls, beam edges, eaves, corners, cracks, crevices, wall voids, unfinished attics, crawl spaces, under appliances, along wall edges, conduits to light fixtures, pillars and other structures such as, but not limited to, fences, awnings, dumpsters, refuse containers and other areas where flies or fly specks are found.

Do not paint surfaces or place boards/cards or rope wicks within the reach of children, pets or livestock.

PESTS KILLED

Pest Group	Including	Application Method		Comments
		Indoors	Outdoors	
Flies	House Fly	Broadcast ¹ Bait Stations Scatter Application to Glue-Boards or Sticky Cards Water-Diluted Product ² (Paint-On or Rope-Wick)	Broadcast Bait Stations	Use bait stations suitable for flies.
	Stable Fly	Bait Stations Scatter Application to Glue-Boards or Sticky Cards		
	Dump Fly Fruit Flies (including Spotted Wing <i>Drosophila</i>) Nuisance Flies (including Black-eyed Fly, Dung Fly ³)	Broadcast ¹ Bait Stations Scatter Application to Glue-Boards or Sticky Cards Water-Diluted Product ² (Paint-On or Rope-Wick)	Broadcast Bait Stations Scatter Application to Glue-Boards or Sticky Cards Water-Diluted Product (Paint-On or Rope-Wick)	
Cockroaches	such as: American Cockroach German Cockroach Oriental Cockroach	Broadcast ¹ Scatter Application to Glue-Boards or Sticky Cards	Broadcast Scatter Application to Glue-Boards or Sticky Cards	

¹ **DO NOT** broadcast indoors except for walkways of penned-animal facilities.

² **DO NOT** USE DILUTED APPLICATIONS (Paint-On or Rope-Wick) for INDOOR RESIDENTIAL SITES.

³ Not approved for use in California.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Do not subject to temperatures below 32 degrees F. Store product in original container only in a location inaccessible to children and pets. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Not for use or storage in or around the home.

Pesticide Disposal: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Non-refillable container. Do not reuse or refill this container. Completely empty container into application equipment. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

For non-emergency information (e.g., current product information), call Central Garden & Pet at 1-800-347-8272.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of CENTRAL GARDEN & PET COMPANY or Seller. To the extent permitted by applicable law, Buyer and User agree to hold CENTRAL GARDEN & PET COMPANY and Seller harmless for any claims relating to such factors.

CENTRAL GARDEN & PET COMPANY warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of the product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or CENTRAL GARDEN & PET COMPANY, and (2) Buyer and User assume the risk of any such use. TO THE EXTENT PERMITTED BY APPLICABLE LAW, CENTRAL GARDEN & PET COMPANY MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall **CENTRAL GARDEN & PET COMPANY** or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF CENTRAL GARDEN & PET COMPANY AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF CENTRAL GARDEN & PET COMPANY OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

CENTRAL GARDEN & PET COMPANY and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of CENTRAL GARDEN & PET COMPANY.

Manufactured for: **Central Garden & Pet Company**, 1501 East Woodfield Road, 200W, Schaumburg, Illinois 60173

NOTE: This specimen label is for informational purposes only. All uses may not be approved in all states.

See product labeling for use directions.

Cyanarox, Starbar and Starbar with design are registered trademarks of Wellmark International. © 2018-2020 Wellmark International.

FR 21-001

Schaumburg, IL
December, 2020