

CAUTION:

- May cause toxic reactions unless drug is evenly mixed in water at dosages indicated and used according to directions.
- Levels of sulfaquinoxaline higher than 0.025% in feed or 0.012% in water for more than 24 to 36 hours may result in reduced growth rate in chickens as a result of reduced feed or water intake.
- For control of disease outbreaks, medication should be initiated as soon as diagnosis is determined.
- Treated animals must actually consume enough medicated water to provide a necessary dosage of approximately 10 to 45 mg/lb/day in chickens and 3.5 to 55mg/lb/day in turkeys and 6 mg/lb/day in cattle and calves, depending on class of animal, ambient temperature, age, and other factors.
- Prolonged administration of sulfaquinoxaline at higher doses may result in depressed feed or water intake, deposition of sulfaquinoxaline crystals in kidney and interference with normal blood clotting.
- Do not mix or administer in galvanized containers.
- The following word is required on this product to comply with the Agricultural Code of California: HAZARDOUS
- CAUSES SKIN AND EYE BURNS. Avoid contact with eyes, skin, or clothing. In case of contact, flush immediately with water for at least 15 minutes; for eye get medical attention.

WARNING:

- Withdraw use of product 10 days before slaughter.
- A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.
- Do not medicate chickens or turkeys producing eggs for human consumption.
- Not for use in lactating dairy cattle.
- DISPOSE OF ANY WASTE OR UNUSED PORTION PROPERLY.
- KEEP ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.
- Store between 20°-25°C (68°-77°F); excursions permitted between 15°-40°C (59°-104°F).

Sul-Q-Nox[®]

31.92% (Sulfaquinoxaline as sodium and potassium salts)

ANTIMICROBIAL

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

For Use in Animal Drinking Water Only

Sulfonamide

KEEP OUT OF REACH OF CHILDREN

KEEP TIGHTLY SEALED WHEN NOT IN USE

Restricted Drug (California) – Use only as directed. Not for human use.

Net Contents: 1 Gallon (128 fl. oz.)

Manufactured for:
Huvepharma, Inc.
525 Westpark Drive, Suite 230
Peachtree City, GA 30269



HUVEPHARMA and Sul-Q-Nox are registered trademarks of Huvepharma EOOD



Approved by FDA under NADA # 006-891
P08-4001SL
Rev. 01-2022

FOR AUTOMATIC DRINKING WATER PROPORTIONERS		
FOR CONTROL OR TREATMENT OF THE FOLLOWING DISEASES WHEN CAUSED BY ONE OR MORE PATHOGENIC ORGANISMS SUSCEPTIBLE TO SULFAQUINOXALINE		
PREPARE MEDICATED DRINKING WATER FRESH DAILY		
INDICATIONS	DOSAGE PER 128 GALLONS OF WATER	TREATMENT
COCCIDIOSIS — CHICKENS: Control of coccidiosis caused by <i>Eimeria tenella</i> , <i>E. necatrix</i> , <i>E. maxima</i> , <i>E. brunetti</i> , <i>E. acervulina</i> , susceptible to sulfaquinoxaline.	20½ fl. oz. (0.04%) 13 fl. oz. (0.025%)	Give for 2-3 days, skip 3 days. — THEN — Give for 2 days more. If bloody droppings appear, repeat treatment at this level for 2 more days. CAUTION: Do not change litter unless absolutely necessary. Do not give flushing mash.
COCCIDIOSIS — TURKEYS: Control of coccidiosis caused by <i>Eimeria meleagrimitis</i> , <i>E. adenoides</i> , susceptible to sulfaquinoxaline.	13 fl. oz. (0.025%)	Give for 2 days, skip 3 days – give for 2 days, skip 3 days and give 2 more days. Repeat if necessary. CAUTION: Do not change litter unless absolutely necessary. Do not give flushing mash.
ACUTE FOWL CHOLERA — CHICKENS AND TURKEYS: Control of acute fowl cholera caused by <i>Pasteurella multocida</i> , susceptible to sulfaquinoxaline.	20½ fl. oz. (0.04%)	Use for 2-3 days. Move birds to clean ground. If disease recurs, repeat treatment. Poultry that have survived fowl typhoid outbreaks should not be kept for laying house replacements or breeders unless tests show they are not carriers.
FOWL TYPHOID — CHICKENS AND TURKEYS: Control of fowl typhoid caused by <i>Salmonella gallinarum</i> , susceptible to sulfaquinoxaline.	20½ fl. oz. (0.04%)	Use for 2-3 days. Move birds to clean ground. If disease recurs, repeat treatment. Poultry that have survived fowl typhoid outbreaks should not be kept for laying house replacements or breeders unless tests show they are not carriers.
COCCIDIOSIS — CATTLE AND CALVES: For the control and treatment of coccidiosis caused by <i>Eimeria bovis</i> <i>E. zuernii</i> , susceptible to sulfaquinoxaline.	7¼ fl. oz. (0.015%) 6 mg/lb of body wt.	Give for 3-5 days. As a generalization, cattle and calves will consume approx. 1 gal/100 lbs body weight/day. Cattle or calves not eating or drinking must be treated individually.
To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Huvepharma, Inc. at 1-877-994-4883 or www.huvepharma.us. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VEIS or http://www.fda.gov/reportanimalae.		

Time/Date: 30066900.pdf 1 12/24/14 3:29 PM

Project No.	Artwork Number	Description	Country
12288	30066900	Sul-Q-Nox	US
Dimensions		Drawing No	SKU No.
18.38" x 5.25"		DWG-300177-00	3800000
Item		Label	
18.38" x 5.25"		Label	
Additional Info: Sleeve Label		Colors: Black Dieline	<input type="checkbox"/> EDITOR'S COPY GS: DATE:
Mgr	Rev	GA	PR
GS	2	GA	PR
GA	2	GA	PR
		CHANGES OK	CHANGES OK
		CHANGES OK	CHANGES OK

Sul-Q-Nox[®]

31.92% (Sulfaquinoxaline as sodium and potassium salts)

ANTIMICROBIAL

For Use in Animal Drinking Water Only

Sulfonamide

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

**KEEP OUT OF REACH OF CHILDREN
KEEP TIGHTLY SEALED WHEN NOT IN USE**

Restricted Drug (California) – Use only as directed. Not for human use.

Net Contents: 4 x 1 Gallon

Approved by FDA under NADA # 006-891

Consult Bottle Label for Use Instructions

Keep sealed tightly when not in use.

Store between 20°-25°C (68°-77°F); excursions permitted between 15°-40°C (59°-104°F).



Manufactured for:
Huvepharma, Inc.
525 Westpark Drive, Suite 230
Peachtree City, GA 30269

HUVEPHARMA and Sul-Q-Nox are registered trademarks of Huvepharma EOOD



DOT SHIPPING INFORMATION

Caustic Alkali Liquid, n.o.s.
(sodium hydroxide,
potassium hydroxide)
UN1719 Class 8 PGII

P08-4001CL
Rev. 01-2022

Time/Date:

zoetis	Project No.	Artwork Number	Description		Country	
	12288	30066800	Sul-Q-Nox		US	
Dimensions		Drawing No	SKU No.	Item		
11" x 7"		NA	3800000	Label		
Additional Info:		Colors: Black PMS 1765 Process Yellow Dieline			<input type="checkbox"/> EDITOR'S COPY GS: DATE:	
Mgr	J. Rustad	Rev	GA	PR	GS / ART REV (LCA)	GS / ART REV (FA)
GS	K Minard/L. Hicks	2			CHANGES OK	CHANGES OK
GA	J. Disch					CHANGES OK