

SulfaMed-G[®] (sulfadimethoxine) Soluble Powder

FOR ORAL USE IN CHICKENS, TURKEYS AND CATTLE

Antibacterial

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian. Federal law prohibits the extralabel use of this product in lactating dairy cattle.

INDICATIONS:

For Broiler and Replacement Chickens Only - Use for the treatment of disease outbreaks of coccidiosis, fowl cholera, and infectious coryza.

For Meat-producing Turkeys Only - Use for the treatment of disease outbreaks of coccidiosis and fowl cholera.

For Dairy Calves, Dairy Heifers, and Beef Cattle - For the treatment of shipping fever complex and bacterial pneumonia associated with *Pasteurella* spp. sensitive to sulfadimethoxine; and calf diphtheria and foot rot associated with *Fusobacterium necrophorum* (*Sphaerophorus necrophorus*) sensitive to sulfadimethoxine.

NOT FOR HUMAN USE

KEEP OUT OF REACH OF CHILDREN

Restricted Drug (California) - Use Only as Directed

ANADA 200-376, Approved by FDA

Each packet contains 3.34 oz (94.6 g) Sulfadimethoxine in the form of the soluble sodium salt and disodium edetate.



Net Weight: 107 g (3.77 oz)

Manufactured By:
Bimeda, Inc.
Le Sueur, MN 56058
www.bimeda.com

**2" gap
bottom of pouch**

Product No.: 1SDM005 8SDM009 Rev. 08/16

MADE IN USA

TAKE TIME

DIRECTIONS

Store at controlled room temperature 20°C - 25°C (68°F - 77°F), with excursions permitted to 15°C - 30°C (59°F - 86°F).

RESIDUE WARNINGS: CHICKENS AND TURKEYS - Withdraw 7 days before slaughter. For dairy calves, dairy heifers and beef cattle only. A withdrawal period has not been established for this product in pre-ruminating calves. Do Not Use in Calves to be Processed for Veal.

PRECAUTIONS: Chickens and turkeys that have survived fowl cholera outbreaks should not be kept for replacements or breeders. Treatment should not be continued beyond 5 days.

Cattle: During treatment period, make certain that animals maintain adequate water intake. If animals show no improvement within 2 or 3 days, re-evaluate diagnosis. dilutions (chickens 0.05% and turkeys 0.025%) as regular drinking water. Administer as sole source of drinking water and sulfonamide medication.

DIRECTIONS FOR USE:

Poultry: Treatment Period 5 consecutive days. Six-600 lb animals initially or twelve-600 lb animals on a maintenance dose. Contents of packet will medicate twenty fl. oz. of cattle stock solution or two-600 lb animal initially or two-600 lb animals on a maintenance dose. Contents of packet will medicate For treatment of individual cattle, SulfaMed-G[®] Soluble Powder stock solution for cattle may be given as a drench. Administer using same mg/lb dosage as compensate for this and insure proper drug intake.

Automatic Proportions - To make stock solution, add contents of 5 packets to 2 gallons of water for chickens and to 4 gallons of water for turkeys. Set proportioner to feed at a rate of 1 fl oz of stock solution per gallon of water.

Treatment Period 6 consecutive days. Dose: 25 mg/lb first day followed by 12.5 mg/lb/day for 4 days.

Species	Concentration	Use Directions
Chickens	0.05%	Contents of packet to 50 gallons of water
Turkeys	0.025%	Contents of packet to 100 gallons of water

SulfaMed-G[®] in Water

Amount of Stock Solution for Cattle*	(Summer) 1 gallon/**100 lb body weight	(Winter) 1 gallon/**150 lb body weight
1 quart	10 gallons	7 gallons
2 quarts	20 gallons	14 gallons
1 gallon	40 gallons	28 gallons
1 quart	20 gallons	14 gallons
2 quarts	40 gallons	28 gallons
1 gallon	80 gallons	56 gallons

Next 4 Days Add

* Note: Make a cattle stock solution by adding 1 packet of SulfaMed-G[®] Soluble Powder to 1 gallon of water. ** This dosage recommendation is based on a water consumption rate of 1 gallon per 100 lb of body weight per day, the expected water consumption rate for summer. Water consumption during cold months (winter) may drop markedly (30-40%). Accordingly, adjustments must be made in the dilution rates to compensate for this and insure proper drug intake.

For treatment of individual cattle, SulfaMed-G[®] Soluble Powder stock solution for cattle may be given as a drench. Administer using same mg/lb dosage as outlined above.

Twenty fl. oz. of cattle stock solution will medicate one-600 lb animal initially or two-600 lb animals on a maintenance dose. Contents of packet will medicate six-600 lb animals initially or twelve-600 lb animals on a maintenance dose.

DIRECTIONS FOR USE:

Poultry: Treatment Period 5 consecutive days.

Cattle: During treatment period, make certain that animals maintain adequate water intake. If animals show no improvement within 2 or 3 days, re-evaluate diagnosis. dilutions (chickens 0.05% and turkeys 0.025%) as regular drinking water. Administer as sole source of drinking water and sulfonamide medication.

RESIDUE WARNINGS: CHICKENS AND TURKEYS - Withdraw 7 days before slaughter. For dairy calves, dairy heifers and beef cattle only. A withdrawal period has not been established for this product in pre-ruminating calves. Do Not Use in Calves to be Processed for Veal.

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian. Federal law prohibits the extralabel use of this product in lactating dairy cattle.