



SulfaMed-G[®] (sulfadimethoxine) Soluble Powder

FOR ORAL USE IN CHICKENS, TURKEYS AND CATTLE

Antibacterial

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian. Federal law prohibits the extralabel use of this product in lactating dairy cattle.

INDICATIONS:

For Broiler and Replacement Chickens Only - Use for the treatment of disease outbreaks of coccidiosis, fowl cholera, and infectious coryza.

For Meat-producing Turkeys Only - Use for the treatment of disease outbreaks of coccidiosis and fowl cholera.

For Dairy Calves, Dairy Heifers, and Beef Cattle - For the treatment of shipping fever complex and bacterial pneumonia associated with *Pasteurella* spp. sensitive to sulfadimethoxine; and calf diphtheria and foot rot associated with *Fusobacterium necrophorum* (*Sphaerophorus necrophorus*) sensitive to sulfadimethoxine.

NOT FOR HUMAN USE

KEEP OUT OF REACH OF CHILDREN

Restricted Drug (California) - Use Only as Directed

ANADA 200-376, Approved by FDA

Each packet contains 16.8 oz (473 g) Sulfadimethoxine in the form of the soluble sodium salt and disodium edetate.



Net
Weight: 535 g (18.87 oz)

Manufactured By:
Bimeda, Inc.
Le Sueur, MN 56058
www.bimeda.com

2.5" gap
bottom of pouch

Product No: 1SDM006 8SDM011 Rev. 08/16



OBSERVE LABEL
DIRECTIONS



TAKE TIME



Store at controlled room temperature 20°C - 25°C (68°F - 77°F), with excursions permitted to 15°C - 30°C (59°F - 86°F).
SulfaMed-G[®] is a Registered Trademark of Bimeda, Inc.
N. A. Corp. Address: Bimeda, Inc., One Tower Lane, Oakbrook Terrace, IL 60181

RESIDUE WARNINGS: CHICKENS AND TURKEYS – Withdraw 5 days before slaughter. Do not administer to chickens over 16 weeks (112 days) of age or to turkeys over 24 weeks (168 days) of age.
CATTLE – Withdraw 7 days before slaughter. For dairy calves, dairy heifers and beef cattle only. A withdrawal period has not been established for this product in pre-ruminating calves. Do Not Use in Calves to be Processed for Veal.
During treatment period, make certain that animals maintain adequate water intake. If animals show no improvement within 2 or 3 days, re-evaluate diagnosis. Treatment should not be continued beyond 5 days.
PRECAUTIONS: Chickens and turkeys that have survived fowl cholera outbreaks should not be kept for replacements or breeders.
Cattle: If animals show no improvement within 5 days, discontinue treatment and re-evaluate diagnosis. Prepare a fresh stock solution daily. Handle the recommended dilutions (chickens 0.05% and turkeys 0.025%) as regular drinking water. Administer as sole source of drinking water and sulfonamide medication.

Directions for Use:
Treatment Period 5 consecutive days.
Twenty fl. oz. of cattle stock solution will medicate one-600 lb animal initially or two-600 lb animals on a maintenance dose. Contents of packet will medicate thirty-600 lb animals initially or sixty-600 lb animals on a maintenance dose.
For treatment of individual cattle, SulfaMed-G[®] Soluble Powder stock solution for cattle may be given as a drench. Administer using same mg/lb dosage as outlined above.
* This dosage recommendation is based on a water consumption rate of 1 gallon per 100 lb of body weight per day, the expected water consumption rate for summer. Water consumption during cold months (winter) may drop markedly (30-40%). Accordingly, adjustments must be made in the dilution rates to compensate for this and insure proper drug intake.
** Note: Make a cattle stock solution by adding 1 packet of SulfaMed-G[®] Soluble Powder to 5 gallons of water.

Species	Concentration	SulfaMed-G [®] in Water	
		Amount of Stock Solution for Cattle*	Water Consumption
Chickens Turkeys	0.025%	1 quart	14 gallons
		2 quarts	28 gallons
Cattle	0.05%	1 quart	14 gallons
		2 quarts	28 gallons
Amount of Stock Solution for Cattle*	1 gallon/100 lb body weight	1 quart	14 gallons
		2 quarts	28 gallons
Next 4 Days Add	1 quart	20 gallons	80 gallons
		2 quarts	40 gallons
First Day Add	1 quart	10 gallons	40 gallons
		2 quarts	20 gallons
Amount of Stock Solution for Cattle*	1 gallon/100 lb body weight	7 gallons	28 gallons
		14 gallons	56 gallons

Automatic Proportions – To make stock solution, add contents of 1 packet to 2 gallons of water for chickens and to 4 gallons of water for turkeys. Set proportioner to feed at a rate of 1 fl. oz of stock solution per gallon of water.
Treatment Period 6 consecutive days. Dosage: 25 mg/lb first day followed by 12.5 mg/lb/day for 4 days.

Species	Concentration	Use Directions
Chickens Turkeys	0.025%	Contents of packet to 500 gallons of water
	0.05%	Contents of packet to 250 gallons of water

DOSE AND ADMINISTRATION:

SulfaMed-G[®] (sulfadimethoxine) Soluble Powder
FOR ORAL USE IN CHICKENS, TURKEYS AND CATTLE