

PRECAUTIONARY STATEMENTS
Hazards to humans and domestic animals

DANGER
KEEP OUT OF REACH OF CHILDREN. Corrosive. Causes irreversible eye damage and skin burns. May be fatal if inhaled. Harmful if swallowed or absorbed through the skin. Do not get in eyes, on skin, or on clothing. Do not breathe spray mist. Wear goggles or face shield. Wear a dust/mist filtering respirator (MSHA/NIOSH approval number TC-21C) or a NIOSH-approved respirator with any N, P, R, or HE pre-filter. Wear protective clothing and rubber gloves. Prolonged or frequently repeated skin contact may cause an allergic reaction in some individuals. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash it before reuse.

ENVIRONMENTAL HAZARDS: This product is toxic to fish, aquatic invertebrates, oysters, and shrimp. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

DESCRIPTION
SYNERGIZE is a multi-purpose disinfectant-cleanser. **SYNERGIZE** provides a broad spectrum kill of gram negative and gram positive microorganisms. **SYNERGIZE** provides effective disinfecting and cleaning in one operation. **SYNERGIZE** is effective in warm, soft, or hard water (as much as 1,000 ppm CaCO₃) and performs especially well in areas containing 5% serum.

SYNERGIZE is for use on glazed ceramic and tile surfaces, stainless steel, aluminum, chrome, galvanized metal, tin, brass, zinc, glass, hard non-porous treated wood, polyethylene, polypropylene, PVC (polyvinyl chloride), vinyl, sealed fiberglass, Viton, ethylene propylene, nitrile, acrylic, and polyurethane.

DIRECTIONS FOR USE:
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Registered for disinfecting and cleaning hard, non-porous surfaces. Not for use on feed, food, or litter.

USE DIRECTIONS:
 To disinfect and clean hard, non-porous surfaces of walls, floors, tables, sinks, and garbage cans in non-animal areas associated with the sites noted on this label. Apply a solution of ½ ounce **SYNERGIZE** per gallon of water (dilute 1:256) to pre-cleaned surfaces. Apply with cloth, sponge, mop, or mechanical coarse spray device, making sure all surfaces are wetted thoroughly. Allow surfaces to remain wet for 10 minutes. All surfaces must be rinsed with clean water and allowed to dry before reuse.

BOOT/SHOE WASH: To prevent cross contamination into animal areas, shoe baths containing one inch of freshly made disinfecting solution must be placed at all entrances to buildings. Soak waterproof shoes or boots and place in a ½ ounce **SYNERGIZE** per gallon of water solution. Allow the solution to come in contact with the boot/shoe for 10 minutes. Change the disinfecting solution when solution becomes visibly dirty.

VEHICLES: To disinfect the non-porous, hard surfaces of vehicles use a solution of ½ ounce **SYNERGIZE** per gallon of water. Apply using a high pressure or coarse spray system. Leave all treated surfaces exposed to solution for 10 minutes or more, and allow to air dry. All treated surfaces that will contact feed or drinking water must be thoroughly scrubbed with soap or detergent then rinsed with potable water before reuse.

- DIRECTIONS FOR USE IN SWINE-PRODUCING FACILITIES:**
- 1) Remove all animals and feed from the premises, vehicles, and enclosures.
 - 2) Empty all troughs, racks, and other feeding and watering appliances.
 - 3) Remove all gross soil, bedding, manure, caked-on dirt, spider webs, etc., from walls to ceilings of barns, pens, stalls, chutes, and other facilities occupied or traversed by hogs or other animals.
 - 4) Thoroughly clean all surfaces with soap or detergent and rinse with water.
 - 5) Thoroughly saturate with a solution of ½ ounce **SYNERGIZE** per gallon of water (1:256), using conventional spraying methods.
 - 6) Allow all treated areas to remain in contact with this solution for at least 10 minutes.
 - 7) Immerse equipment used in handling and restraining animals, forks, shovels, and scrapers used for removing manure in **SYNERGIZE** at a dilution of 1:256. Allow to remain in contact for at least 10 minutes.
 - 8) Ventilate buildings and other closed spaces. Do not house swine or other livestock until treatment has been absorbed, set, or dried.
 - 9) Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains, waterers, and equipment which may come in contact with food with soap or detergent, and rinse with potable water before reuse.

- DIRECTIONS FOR USE IN FARM PREMISES, EQUINE FACILITIES, VEAL BARN, RATITES FACILITIES, CALF PENS, AND VEHICLES:**
- 1) Do not use in milking stalls, milking parlors, or milk houses.
 - 2) Remove all animals and feed from the premises, vehicles, and enclosures.
 - 3) Remove all gross soil, bedding, manure, caked-on dirt, spider webs, etc., from floors, surfaces, and from walls to ceiling of barns, pens, stalls, chutes, and other facilities occupied or traversed by animals.
 - 4) Empty all troughs, racks, and other feeding and watering appliances.
 - 5) Thoroughly clean all surfaces with soap or detergent and rinse with water.
 - 6) Thoroughly saturate with a solution of ½ ounce **SYNERGIZE** per gallon of water (1:256), using conventional spraying methods. Allow to remain in contact with this solution for at least 10 minutes.
 - 7) Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing manure in a solution of ½ ounce **SYNERGIZE** per gallon of water (1:256). Allow to remain in contact for 10 minutes.
 - 8) Ventilate buildings, cars, boats, and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set, or dried.
 - 9) Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains, waterers, and equipment which may come in contact with food with soap or detergent, and rinse with potable water before reuse.

- DIRECTIONS FOR USE IN POULTRY HOUSES:**
- 1) Remove all poultry and feeds from premises, trucks, coops, and crates.
 - 2) Remove all litter and droppings from floors, walls, and surfaces of facilities occupied or traversed by poultry.
 - 3) Empty all troughs, racks, and other feeding and watering appliances.
 - 4) Thoroughly clean all surfaces with soap or detergent and rinse with water.
 - 5) Saturate surfaces with ½ ounce **SYNERGIZE** per gallon of water (dilute 1:256) for a period of 10 minutes.
 - 6) Ventilate buildings, coops, and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set, or dried.
 - 7) Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains, waterers, and equipment which may come in contact with food with soap or detergent, and rinse with potable water before reuse.

HATCHERY ROOMS, INCUBATORS AND HATCHERS, POULTRY HOUSES, AND LIVESTOCK BUILDINGS BY FOGGING:
 All surfaces must be cleaned and disinfected in accordance with label directions prior to fogging. Fogging is not to be used as a sanitizing treatment. Close room off so fog is confined to room to be treated. Hatchery ventilation systems must remain on during fogging. Wear a dust/mist respirator when mixing the use solution and pouring it into the fogging apparatus.

SYNERGIZE™

CLEANER • DISINFECTANT

STAPHYLOCIDAL • PSEUDOMONACIDAL • SALMONELLACIDAL • VIRUCIDAL • FUNGICIDAL • ASPERGILLICIDAL

Synergize cleaner-disinfectant is specifically formulated for industrial use in the following Animal and Poultry Housing facilities: Turkey, Breeder, Grow Out, Fryer, Layer, Hatcheries and Swine-Producing Facilities, as well as the following equipment in these facilities: Waterers, Feeders, Hauling Equipment, Loading Equipment, Farrowing Bins, Nurseries, Blocks, Creep Areas, and Chutes. This product is also for use in Farm Premises, Equine Facilities, Veal Barns, Ratite Facilities (Ostrich, Emu and Rhea Facilities), Calf Pens, and associated Farm Vehicles.

Active Ingredients:

Alkyl (C₁₂ 67%, C₁₄ 25%, C₁₆ 7%, C₁₈ 1%)
 dimethyl benzyl ammonium chloride.....26.00%
 Glutaraldehyde.....7.00%

Inert Ingredients:.....67.00%
 TOTAL.....100.00%

EPA Reg. No. 66171-7
 Patent Number 5,891,922

EPA Est. No. 66171-TN-001
 EPA Est. No. 66171-CA-001
 EPA Est. No. 61282-WI-01
 EPA Est. No. 47000-IA-01

(First letter of batch number indicates establishment)

KEEP OUT OF REACH OF CHILDREN DANGER

See side panels for additional Precautionary Statements

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor or going in for treatment. For 24-hour emergency information on this product, call 1-800-498-5743 (US & Canada) or 1-651-523-0318 (all other areas).

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsions may be needed.



Net Contents: 55 gal. (208 L)

Directions for Use, continued

A) HATCHERY ROOMS:
 Vacate all personnel from hatchery room to be fogged. Close room off so fog is confined to the hatchery room to be treated. Prepare a stock solution of one (1) part **SYNERGIZE** to forty-two (42) parts water (3 fluid ounces **SYNERGIZE** to 125 fluid ounces of water). Fog this solution at a rate of 3.8 ounces per 10,000 cubic feet for the fogging times specified in the chart below. Contact your Preserve International representative to determine the recommended fogging schedule. Do not allow people to breathe or contact the fog. When fogging **SYNERGIZE**, re-entry intervals (REI) and personal protective equipment (PPE) are determined by the following hatchery room air exchange rates:

ACH for Hatchery Rooms/Incubators and Hatchers	Fogging Time	Re-entry Intervals (REI)	PPE Required
8 Air Exchanges Per Hour	5 Minutes	25 Minutes	No PPE Required
12 Air Exchanges Per Hour	7.5 Minutes	15 Minutes	No PPE Required
16 Air Exchanges Per Hour	10 Minutes	10 Minutes	No PPE Required
20 Air Exchanges Per Hour	12.5 Minutes	10 Minutes	No PPE Required
24 Air Exchanges Per Hour	15 Minutes	10 Minutes	No PPE Required

Under no circumstances must the hatchery room be entered by unprotected persons before the listed re-entry intervals. If the hatchery room must be entered before the listed re-entry intervals, then the individual entering the room must wear a long-sleeved shirt and pants and an approved NIOSH/MSHA full face piece respirator, such as the 3M 6700-6900 series, with an organic vapor cartridge, such as the 3M 6001, and a particulate filter, such as the 3M 5N11 & 3M 501 filter retainer.

B) INCUBATORS AND HATCHERS:
 Only for treatment of setters and hatchers after poultry/chicks/eggs have been removed. Not for treatment of hatchers which contain chicks/eggs. Prepare a stock solution of one (1) part **SYNERGIZE** to forty-two (42) parts water (3 fluid ounces **SYNERGIZE** to 125 fluid ounces of water). Fog this solution at a rate of 3.8 ounces per 10,000 per cubic feet for the fogging times specified in the chart above, into setters and hatchers immediately after transfer. Repeat daily. Discontinue hatchery treatments approximately 24 hours before pulling the hatch. Do not allow people to contact or breathe this fog and do not enter machines until the fog has been exhausted. To do this, install permanent fogging nozzles in setters and hatchers and use an air compressor to disperse the sanitizing solution as a fog. It is also satisfactory to fog setters and hatchers with a 1:1,000 solution of **SYNERGIZE**. If this is done, fog for 30 – 90 seconds once per hour or once every two hours.

C) POULTRY HOUSES AND LIVESTOCK BUILDINGS:
 After the house has been depopulated and cleaned as in "Directions for use in farm premises, equine facilities, veal barns, ratite facilities, calf pens, and vehicles" double-check to be sure all people, poultry, livestock, and pets have vacated the building. Close all windows, doors, curtains, etc., making the house as closed as possible. Prepare a stock solution of one (1) part **SYNERGIZE** to four (4) parts water (25 fluid ounces **SYNERGIZE** to 100 fluid ounces of water). Insert the nozzle of the fogging device through a suitable opening in the room. With the setting on maximum output, fog 125 fluid ounces for each 1,000 cubic yards. The fogger itself is placed just inside the door of the building to be treated, or the nozzle of the fogger is inserted through a suitable opening in the door or building. The opening must be large enough to accommodate the nozzle. After fogging, the building must be kept closed for twenty-four (24) hours. After twenty-four (24) hours, the fog must have settled and the house can now be opened and aired. The house must be opened for a minimum of twenty-four (24) hours before it is repopulated with poultry or livestock.

NOTE FOR POULTRY HOUSES AND LIVESTOCK BUILDINGS: The generated fog is very irritating to eyes, skin, and mucous membranes. Under no circumstances must a room or building be entered by anyone until the fog has completely settled. A re-entry time of two (2) hours is required. If the building or room must be entered, then the individual entering the building or room must wear a self-contained respirator approved by NIOSH/MSHA, goggles, long-sleeved shirt, and pants. If feeders and waterers were not removed from the premises during treatment, or were not adequately covered to prevent contact with the treatment, they must be washed with detergent and water before use for poultry or livestock.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE:
 Store away from heat. Do not transfer to another container. Do not reuse empty container

PESTICIDE DISPOSAL:
 Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available, or offer for reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Dispose of rinsate as noted in the pesticide disposal section. Repeat this procedure two more times.

General: Consult Federal, State, or local disposal authorities for approved alternative procedures.

LIMITED WARRANTY

To the extent permitted by applicable law, Neogen Corporation makes no warranty concerning uses which extend beyond the use of the product under normal conditions in accordance with the statements made on this label. To the extent permitted by applicable law, Neogen Corporation shall not be liable for (1) any consequential, incidental, or special damages related in any way to this product or its uses, or (2) any damages related in any way to resistance to pesticides.

For non-emergency (e.g., current product information), call: 1-800-621-8829.

Item No.: 433600/55



Manufactured by: Preserve International
 944 Mandino Blvd., Lexington, KY 40511 U.S.A.
 Preserve International is a wholly-owned subsidiary of
 Neogen Corporation.
 L4772-1017

