

Terra-Vet™ 200 (200 mg/mL) (oxytetracycline injection) is a sterile preconstituted solution of the broad spectrum antibiotic base oxytetracycline. Each mL contains 200 mg of oxytetracycline base as oxytetracycline amphoteric and on a w/w basis, 40.0% 2-pyrrolidone, 5.0% povidone, 1.8% magnesium oxide, 0.2% sodium formaldehyde sulfoxylate (as a preservative), monoethanolamine and/or hydrochloric acid as required to adjust pH.

**WARNING:**

Discontinue treatment at least 28 days prior to slaughter of cattle and swine. Milk taken from animals during treatment and for 96 hours after the last treatment must not be used for food.

**PRECAUTIONS:**

Exceeding the highest recommended level of drug per pound of body weight per day, administering more than the recommended number of treatments, and/or exceeding 10 mL intramuscularly or subcutaneously per injection site in adult beef and dairy cattle, and 5 mL intramuscularly per injection site in adult swine, may result in antibiotic residues beyond the withdrawal period.

Made in UK.

Lot No:

8201-92

Exp Date:

MAY-2010



ANADA 200-306, APPROVED BY FDA

**Terra-Vet™ 200**

**(oxytetracycline injection)**

**(200 mg/mL)**

**ANTIBIOTIC**

Each mL contains 200 mg of oxytetracycline base as amphoteric oxytetracycline

For the treatment of disease in beef cattle; dairy cattle; calves, including preruminating (veal) calves; and swine



**NET CONTENTS:**  
**500 mL**

**DOSAGE:**

Terra-Vet™ 200 (200 mg/mL)

**CATTLE:**

A single dosage of 9 milligrams of oxytetracycline per pound of body weight (4.5 mL/100 lb) administered intramuscularly or subcutaneously is recommended in the treatment of the following conditions:

- (1) bacterial pneumonia caused by *Pasteurella* spp. (shipping fever) in calves and yearlings, where retreatment is impractical due to husbandry conditions, such as cattle on range, or where repeated restraint is inadvisable.
- (2) infectious bovine keratoconjunctivitis (pink eye) caused by *Moraxella bovis*.

**SWINE:**

A single dose of 9 milligrams of oxytetracycline per pound of body weight (4.5 mL/100 lb) administered *intramuscularly* is recommended in the treatment of bacterial pneumonia caused by *Pasteurella multocida* in swine, where retreatment is impractical due to husbandry conditions or where repeated restraint is inadvisable.

Refer to package insert for complete indications, dosage and usage. Store at room temperature 15° - 30°C (59° - 86°F). Keep from freezing.

Manufactured for:

Aspen Veterinary Resources,® Ltd.  
Liberty, MO 64068, USA

02162BL03

