



Terramycin® 200

(oxytetracycline)



FOR USE IN DRY FEEDS ONLY. NOT FOR USE IN LIQUID FEED SUPPLEMENTS.

Mixing and Use Directions:
Thoroughly mix the amount of this dry feed with an equal amount by weight of feed for each ingredient prior to blending into a complete feed.

Indications for Use	Oxytetracycline Amount	lb. of Terramycin 200 lbs.
CHICKENS		
Control of infectious organisms caused by <i>Mycoplasma gallisepticum</i> ; control of bacterial diseases caused by <i>Chlamydia coli</i> or <i>Chlamydia psittaci</i> in susceptible to oxytetracycline	600 gms per ton Feed continuously for 7-14 days	0.5-1
Control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Chlamydia coli</i> susceptible to oxytetracycline	400 gms per ton Feed continuously for 7-14 days	2
Reduction of mortality due to air sacculitis (air sac infection) caused by <i>Chlamydia coli</i> susceptible to oxytetracycline	500 gms per ton Feed continuously for 7 days	2.5
<p>⚠ WARNING: Do not use in low calcium feed containing less than 0.50% dietary calcium. Use in such feeds may result in a relative resistance. Do not administer to chickens producing eggs for human consumption.</p>		
TURKEYS		
Control of bacteremia caused by <i>Salmonella mbandaka</i> susceptible to oxytetracycline	100 gms per ton Feed continuously for 7-14 days	0.5
Control of infectious organisms caused by <i>Mycoplasma gallisepticum</i> susceptible to oxytetracycline	200 gms per ton Feed continuously for 7-14 days	1
Control of purulent bacterial organisms associated with <i>Chlamydia psittaci</i> (respiratory tract), <i>Chlamydia coli</i> (air sacs), susceptible to oxytetracycline	25 mg/lb of body weight daily Feed continuously for 7-14 days	4.10*
<p>⚠ WARNING: Do not use in low calcium feed or higher, withdraw 5 days before slaughter. Do not administer to turkeys producing eggs for human consumption.</p>		
SWINE		
Treatment of bacterial enteritis caused by <i>Salmonella choleraesuis</i> and <i>Shigella flexneri</i> or susceptible to dry therapy class and treatment of bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline	10 mg/lb of body weight daily Feed continuously for 7-14 days	2.5*
For breeding swine for control and treatment of <i>Leptospira</i> (reducing the incidence of abortion and stillbirth of offspring) caused by <i>Leptospira pomona</i> susceptible to oxytetracycline	10 mg/lb of body weight daily Feed continuously for two weeks after 14 days	2.5*
CALVES, BEEF CATTLE, AND MONA CASTRING ON DRY CATTLE		
For growing cattle (up to 400 lb) for the reduction of the incidence of bacterial enteritis or susceptible to dry therapy class and treatment of bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline	75 mg/head/day Feed continuously	0.375*
Prevention and treatment of the early stages of shipping fever complex. (Feed 3-5 days before and after arrival at feedlot.)	0.5-2.0 g/head/day	2.5-10*
Treatment of bacterial enteritis caused by <i>Salmonella choleraesuis</i> and bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline	10 mg/lb of body weight daily Feed continuously for 7-14 days	20*
SWEEP		
Treatment of bacterial enteritis caused by <i>Salmonella choleraesuis</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline	10 mg/lb of body weight daily Feed continuously for 7-14 days	0*

⚠ **WARNING:** 5-day withdrawal before slaughter.

* If bird weighs 10 lb, consuming 0.5 lb of complete feed per day
 * If pig consuming 4 lb of complete feed per 100 lb body weight per day
 * Include in feed supplements based on consumption of 2 lb of supplements per head per day
 * If animal weighs 500 lb, consuming 2 lb of supplements per head per day
 * If lamb weighs 60 lb, consuming 1 lb of supplements per head per day

RESTRICTED DRUG (CALIFORNIA) - USE ONLY AS DIRECTED
 NOT FOR HUMAN USE

Developed by:
 Phibro Animal Health Corporation
 Teaneck, NJ 07666

Terramycin® 200

(oxytetracycline)



Terramycin® 200

(oxytetracycline)

TYPE A
 MEDICATED ARTICLE

Active Drug Ingredient:
 Oxytetracycline (from oxytetracycline dihydrate base) equivalent to oxytetracycline hydrochloride 200 g/lb

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

CAUTION: For use in manufacturing medicated animal feeds only.

CAUTION: Certain components of animal feeds, including medicated premixes, possess properties that may be a potential health hazard or a source of personal discomfort to certain individuals who are exposed to them. Human exposure should, therefore, be minimized by observing the general industry standards for occupational health and safety.

Precautions such as the following should be considered: dust masks or respirators and protective clothing should be worn; dust-arresting equipment and adequate ventilation should be utilized; personal hygiene should be observed; wash before eating or leaving a work site; be alert for signs of allergic reactions—seek prompt medical treatment if such reactions are suspected.

STORE AT OR BELOW 25°C (77°F), EXCURSIONS PERMITTED UP TO 40°C (104°F)

SEE BACK PANEL FOR COMPLETE MIXING DIRECTIONS, USE DIRECTIONS, AND WARNINGS

TERRAMYCIN is a registered trademark of Pfizer, Inc licensed to Phibro Animal Health Corporation for Oxytetracycline PHIBRO and PHIBRO logo design are trademarks owned by Phibro Animal Health Corporation or its affiliates. Phibro Animal Health Corporation, Teaneck, NJ 07666

Net Weight 50 lb (22.7 kg)

