



INDICATIONS: For use in the control and treatment of the following conditions in swine, calves and poultry.

FOR SWINE AND CALVES

INDICATIONS: Use for the control and treatment of bacterial enteritis (scours) caused by *Escherichia coli*; bacterial pneumonia associated with *Actinobacillus pleuropneumoniae*, *Pasteurella* spp., and *Klebsiella* spp. sensitive to tetracycline hydrochloride.

RECOMMENDED DOSAGE LEVEL: Use soluble powder in the drinking water at a drug level of tetracycline hydrochloride per gallon to provide 10 mg/lb of body weight per day in divided doses.

RESIDUE WARNING: SWINE - Do not slaughter animals for food purposes within 4 days of treatment.

CALVES - Do not slaughter animals for food purposes within 5 days of treatment. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

CAUTION: Do not use for more than 5 consecutive days.

DIRECTIONS FOR USE: Administer for 3-5 days.

FOR CHICKENS

INDICATIONS: Use for the control of chronic respiratory disease (CRD) and air sac disease caused by *Mycoplasma gallisepticum* and *Escherichia coli*; infectious synovitis caused by *Mycoplasma synoviae* sensitive to tetracycline hydrochloride.

RECOMMENDED DOSAGE LEVEL: CRD and air sac disease: Use soluble powder in the drinking water at a drug level of 400-800 mg tetracycline hydrochloride per gallon. Infectious synovitis: Use soluble powder in the drinking water at a drug level of 200-400 mg tetracycline hydrochloride per gallon.

FOR TURKEYS

INDICATIONS: Use for the control of infectious synovitis caused by *Mycoplasma synoviae*; bluecomb (transmissible enteritis, coronaviral enteritis) complicated by organisms sensitive to tetracycline hydrochloride.

RECOMMENDED DOSAGE LEVEL: Infectious synovitis: Use soluble powder in the drinking water at a drug level of 400 mg tetracycline hydrochloride per gallon. Bluecomb: Use soluble powder in the drinking water at a drug level of tetracycline hydrochloride per gallon to provide 25 mg/lb of body weight per day in divided doses.

RESIDUE WARNING: Do not slaughter birds for food within 4 days of treatment.

CAUTION: Not for use in turkeys or chickens producing eggs for human consumption.

Do not use for more than 14 consecutive days.

DIRECTIONS FOR USE: Administer for 7-14 days. Medicate at first clinical signs of disease or when experience indicates the disease may be a problem.

Lot:
Exp:

TetraMed® 324 HCA (tetracycline hydrochloride) Soluble Powder

FOR ORAL USE IN SWINE, CALVES, CHICKENS, AND TURKEYS


Antibiotic

**CAUTION: Federal law restricts this drug to use
by or on the order of a licensed veterinarian.**

**NOT FOR USE IN HUMANS
KEEP OUT OF REACH OF CHILDREN**

**Restricted Drug (California) - Use Only as Directed
ANADA 200-374, Approved by FDA**

Each pound contains 324 g of tetracycline hydrochloride.



**Net Weight:
2.27 kg (5 lb.)**

Manufactured by:
Bimeda, Inc.
Le Sueur, MN 56058
www.bimeda.com

GENERAL CAUTION: When used in plastic or stainless steel waterers prepare fresh solution every 24 hours. When used in galvanized waterers or automatic medicators, prepare fresh solution every 12 hours. Prepare fresh stock solution every 12 hours. Use as sole source of tetracycline. Diagnosis should be reconsidered if improvement is not noticed within 3 days. The concentration of drug required in medicated water must be adequate to compensate for variations in the age and class of animals, feed consumption, and environmental temperature and humidity, each of which affects water consumption.

MIXING DIRECTIONS:

FOR SWINE, CALVES, AND TURKEYS: 2.52 oz. (two scoops) dissolved in 500 mL (approximately 17 fl. oz.) of warm water will provide a stock solution of 100 mg of tetracycline hydrochloride activity per mL.

FOR TURKEYS ONLY: This stock solution when metered at approximately 1 oz. per gallon will provide drinking water containing 2,957 mg of tetracycline hydrochloride activity per gallon.

The contents of two scoops are sufficient to deliver the recommended daily dosage levels as follows:

At 25 mg/lb of body weight.....2,040 total lbs. of turkeys to be medicated.

At 10 mg/lb of body weight.....5,100 total lbs. of swine or calves to be medicated.

Individual calf treatment.....5 mL (1 measuring teaspoonful) twice daily for each 100 lbs. of body weight administered as a drench or by dose syringe.

FOR CHICKENS AND TURKEYS: To arrive at the recommended dosages, use the contents of 2 scoops to prepare stock solutions as follows:

200 mg/gallon - dissolve 2 scoops (2.52 oz.) in 7,570 mL warm water (2 gallons)

400 mg/gallon - dissolve 2 scoops (2.52 oz.) in 3,785 mL warm water (1 gallon)

800 mg/gallon - dissolve 2 scoops (2.52 oz.) in 1,892 mL warm water (0.5 gallon)

These stock solutions should then be metered into drinking water at approximately 1 oz. per gallon.

At 200-800 mg/gallon two scoops will provide 63.8 to 255 gallons of medicated drinking water.

SPECIAL DIRECTIONS FOR BABY CALVES AND BABY PIGS:

Administer this product one hour before or two hours after feeding milk or milk replacers. Provide clean (unmedicated) drinking water at all times.

NOTE: When using a watering trough, this product is to be administered twice a day in the drinking water of swine, calves and poultry. One-half of the recommended daily dosage level of antibiotic is to be consumed during each administration period, thus providing the drug in divided doses. The contents of this container (5 lb) will provide sufficient drug to treat 162,000 total pounds of swine or calves for a single day at the recommended dosage level of 10 mg/lb of body weight in divided doses. Or the contents of this container will treat 64,900 lbs of turkeys when supplied at 25 mg/lb or at 200-800 mg/gallon will provide 2,025 to 8,100 gallons of medicated drinking water for chickens or turkeys.

Store at controlled room temperature 20°C - 25°C (68°F - 77°F), with excursions permitted to 15°C - 30°C (59°F - 86°F).

TetraMed® is a Registered Trademark of Bimeda, Inc.

N.A. Corp. Address: Bimeda, Inc., One Tower Lane, Oakbrook Terrace, IL 60181

Product No. 1TET023 8TET023 Rev. 09/16

