



Tetroxy[®] HCA-1400

(oxytetracycline hydrochloride)
Soluble Powder

**FOR CONTROL AND TREATMENT OF SPECIFIC DISEASES
IN CHICKENS, TURKEYS, AND SWINE**

Antibiotic

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

This packet contains 512 grams of oxytetracycline hydrochloride and will make:

2560 gallons (9690 L) containing 200 mg oxytetracycline hydrochloride per gallon.

1280 gallons (4845 L) containing 400 mg oxytetracycline hydrochloride per gallon.

640 gallons (2422 L) containing 800 mg oxytetracycline hydrochloride per gallon.

This packet will treat 51,200 lbs. of swine.

This packet will treat 20,480 lbs of turkey as 25 mg/lb body weight.

**FOR USE IN DRINKING WATER ONLY
NOT FOR USE IN LIQUID MEDICATED FEEDS
SEE BACK FOR INSTRUCTIONS**

**NOT FOR USE IN HUMANS
KEEP OUT OF REACH OF CHILDREN
HAZARDOUS REMEDY**

Restricted Drug (California) - Use Only As Directed
ANADA 200-144, Approved by FDA



**Net Weight:
3.09 lb. (1400 g)**



Manufactured By:
Bimeda, Inc.
Le Sueur, MN 56058
www.bimeda.com

Tetroxy[®] HCA-1400 (oxytetracycline hydrochloride) Soluble Powder

MIXING INSTRUCTIONS FOR WATER MEDICATION: CHICKENS - TURKEYS - SWINE

DOSAGE AND INDICATIONS FOR USE IN WATER PROPORTIONERS: Add one packet to the following amount of water to make a stock solution for use in a proportioner set to meter at the rate of one ounce per gallon.

	DISEASE	TREATMENT LEVEL	GALLONS STOCK/PACKET
CHICKENS	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	200-400 mg	20-10
	Control of chronic respiratory disease (CRD) and air sac infections caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> , susceptible to oxytetracycline.	400-800 mg	10-5
	Control of fowl cholera caused by <i>Pasteurella multocida</i> susceptible to oxytetracycline.	400-800 mg	10-5
TURKEYS	Control of hexamitiasis caused by <i>Hexamita meleagridis</i> , susceptible to oxytetracycline.	200-400 mg	20-10
	Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> , susceptible to oxytetracycline.	400 mg	10
	GROWING TURKEYS - Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis), susceptible to oxytetracycline.	25 mg/lb body weight	Varies with age & water consumption
SWINE	FOR THE CONTROL AND TREATMENT OF THE FOLLOWING DISEASES IN SWINE: Bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> , susceptible to oxytetracycline. Bacterial pneumonia caused by <i>Pasteurella multocida</i> , susceptible to oxytetracycline. FOR BREEDING SWINE: Leptospirosis (reducing the incidence of abortions and shedding of leptospira) caused by <i>Leptospira pomona</i> , susceptible to oxytetracycline	10 mg/lb body weight	Varies with age & water consumption

DIRECTIONS FOR USE: Mix fresh solutions daily. Use as sole source of drinking water-do not mix this product directly with milk or milk replacers. Administer 1 hour before or 2 hours after feeding milk or milk replacers. Consult a poultry diagnostic laboratory or poultry pathologist for diagnosis and advice on dosage for chickens and turkeys. As a generalization, 200 chickens will drink one gallon of water per day for each week of age. Turkeys will consume twice that amount. Administer 7 to 14 days to chickens & turkeys and up to 5 days to swine.

NOTE: The concentration of drug required in medicated water must be adequate to compensate for variation in the age of the animal, feed consumption rate, and the environmental temperature and humidity, each of which affects water consumption.

PRECAUTION: Use as sole source of oxytetracycline. Not to be used for more than 14 consecutive days in chickens and turkeys or 5 consecutive days in swine.

RESIDUE WARNING:

Do not feed to birds producing eggs for human consumption. Discontinue treatment of chickens, turkeys and swine 0 days prior to slaughter.

RECOMMENDED PACKET STORAGE CONDITIONS: Store between 20°C - 25°C (68°F - 77°F) with excursions permitted between 15°C - 40°C (59°F - 104°F).

CONSULT SAFETY DATA SHEET PRIOR TO USE.



TAKE TIME



OBSERVE LABEL DIRECTIONS



Tetroxy[®] is a registered trademark of Bimeda, Inc.
N.A. Corp. Address: Bimeda, Inc., One Tower Lane, Oakbrook Terrace, IL 60181

Product No. 1TET014
8TET006 Rev. 10/16