

Tetroxy®-LA (oxytetracycline injection) is a sterile solution of the broad-spectrum antibiotic oxytetracycline.

EACH mL CONTAINS: 200 mg of oxytetracycline base as oxytetracycline dihydrate, 2.5% povidone, 1.8% magnesium oxide, 0.3% sodium formaldehyde sulfoxylate, and an aqueous/organic solvent system. Monoethanolamine and/or hydrochloric acid as required to adjust pH.

DOSAGE:

CATTLE- A single dosage of 9 milligrams of oxytetracycline per pound of body weight (4.5 mL/100 lb) administered *intramuscularly* is recommended in the treatment of the following conditions: 1) bacterial pneumonia caused by *Pasteurella* spp. (shipping fever) in calves and yearlings, where re-treatment is impractical due to husbandry conditions, such as cattle on range, or where repeated restraint is inadvisable; 2) infectious bovine keratoconjunctivitis (pinkeye) caused by *Moraxella bovis*.

SWINE- A single dosage of 9 milligrams of oxytetracycline per pound of body weight (4.5 mL/100 lb) administered *intramuscularly* is recommended in the treatment of bacterial pneumonia caused by *Pasteurella multocida* in swine, where re-treatment is impractical due to husbandry conditions or where repeated restraint is inadvisable.

Refer to package insert for complete indications, dosage and usage.



TETROXY[®]-LA

(oxytetracycline injection)

ANTIBIOTIC

FOR ANIMAL USE ONLY
KEEP OUT OF REACH OF CHILDREN

For Use in Beef Cattle, Non-lactating Dairy Cattle, Calves,
Including Pre-ruminating (Veal) Calves, and Swine.

Each mL contains 200 mg of oxytetracycline
base as oxytetracycline dihydrate.

ANADA 200-117, Approved by FDA

Net Contents: 250 mL

Manufactured for:
Bimeda, Inc.
Oakbrook Terrace, IL 60181


Bimeda

WARNING: Discontinue treatment at least 28 days prior to slaughter of cattle and swine. Not for use in lactating dairy animals.

PRECAUTIONS: Exceeding the highest recommended dosage level of drug per pound of body weight per day, administering more than the recommended number of treatments, and/or exceeding 10 mL intramuscularly per injection site in adult beef cattle and non-lactating dairy cattle, and 5 mL intramuscularly per injection site in adult swine, may result in antibiotic residues beyond the withdrawal period.

STORE AT ROOM TEMPERATURE, 15°C - 30°C (59°F - 86°F).

KEEP FROM FREEZING.


Restricted drug (California), use only as directed.

© Registered Trademark of Bimeda, Inc.

Manufactured by: Bimeda-MTC Animal Health Inc.
Cambridge, Ontario, Canada N3C 2W4

Bimeda, Inc. and Bimeda-MTC Animal Health Inc.
are Divisions of Cross Vetpharm Group Ltd.

Product No. 1TET029
8TET023



8