


PRODUCT INFO

Product Name	Tiamulin Hydrogen Fumarate	Product Type	Medicated Swine Feed
Lot No.	1611000	Manufacturer	Denagard, Inc.
Exp. Date	12/31/2018	Product Code	1611000
Product Size	35 lb	Product Weight	15.9 kg

ATTENTION INFO

Product Name	Tiamulin Hydrogen Fumarate	Product Code	1611000
Lot No.	1611000	Manufacturer	Denagard, Inc.
Exp. Date	12/31/2018	Product Weight	15.9 kg
Product Size	35 lb	Product Weight	15.9 kg

LEGEND KEY Tiamulin, BS, Hydrogen Fumarate, CT



tiamulin hydrogen fumarate
Type B Medicated Swine Feed
For manufacture of Type C medicated swine feed only
ACTIVE INGREDIENT: Tiamulin hydrogen fumarate:10 g/lb
Ingredients: rice hulls, mineral oil, porcine gelatin
See reverse side for use directions, cautions and warnings.
Net Weight: 35 lb (15.9 kg)

COVERS
 FRICTION RESISTING BAG
 Meets requirements of
 applicable Federal Specification
 approved by
 military forces
 Bag Size 16x5x34

tiamulin hydrogen fumarate
Type B Medicated Swine Feed
For Manufacture of Type C medicated swine feed only. Do not feed undiluted.

Indications	Amount of Type B Medicated Feed Per Ton of Feed	Tiamulin hydrogen fumarate in Complete Type C Medicated Feed Per Ton	Withdrawal Period (days)
For treatment of swine dysentery associated with <i>Brachyspira</i> (formerly <i>Serpulina</i> or <i>Treponema</i>) <i>hyodysenteriae</i> susceptible to Tiamulin	20 lb	200 g	7
For control of porcine proliferative enteropathies (ileitis) associated with <i>Lawsonia intracellularis</i>	3.5 lb	35 g	2
For control of swine dysentery associated with <i>Brachyspira</i> (formerly <i>Serpulina</i> or <i>Treponema</i>) <i>hyodysenteriae</i> susceptible to Tiamulin	3.5 lb	35 g	2

Warnings:
 Observe withdrawal time shown in table above.
 Keep out of reach of children. Avoid contact with skin. Direct contact with skin or mucous membranes may cause irritation.
Caution:
 Do not feed undiluted. Do not use in feeds for animals other than swine. The effects of tiamulin on swine reproductive performance, pregnancy and lactation have not been determined. Swine being treated with Denagard (tiamulin) should not have access to feeds containing polyether ionophores (e.g., lasalocid, monensin, narasin, salinomycin and semduramicin) as adverse reactions may occur. If signs of toxicity occur, discontinue use.

MIXING DIRECTIONS
 Mix this medicated feed (Type B) with non-medicated feed ingredients to make a Type C medicated feed as follows:

Type C Medicated Feed Tiamulin grams/ton Feed	Pounds Medicated Feed (Type B)	Pounds Non-medicated Feed*
35	3.5	1996.5
200	20	1980

*When making an approved Type C medicated feed drug combination, the inclusion rate of this non-medicated feed should be adjusted for the amount of the other drug added to make such combination Type C medicated feed.

Storage:
 Store at or below 25°C (77°F). Temporary excursions to 40°C (104°F) permitted.
 To report adverse effects, access medical information, or obtain additional product information call 1-800-428-4441
 For a complete listing of adverse reactions for tiamulin hydrogen fumarate reported to CVR see: <http://www.fda.gov/AnimalVeterinary/SafetyHealth>.

Restricted Drug (California): Use only as directed.
 NADA 139-472, Approved by FDA.
 Marketed by:



BC101596X
 PO# XXXXXX
 1681
 Bag Size 16x5x34