



VICTUS FINISHER



Contains a source of endo-beta-glucanase derived from *Trichoderma longibrachiatum*. Endo-beta-glucanase hydrolyzes beta-glucans, a type of non-starch polysaccharide to reduce digesta viscosity with barley based poultry diets and reduces soluble non-starch polysaccharides in digesta. Contains a source of endo-xylanase derived from *Trichoderma longibrachiatum*. Xylanase hydrolyzes xylans, a component of hemicellulose to reduce digesta viscosity with poultry diets. Contains a source of phytase derived from *Aspergillus oryzae* modified by a phytase gene from *Citrobacter braakii*. Phytase increases the digestibility of phytin-bound phosphorus in poultry and swine diets.

GUARANTEED ANALYSIS

Endo-1,4-B-xylanase (T. longibrachiatum), U	Minimum	408 240	900 000
Phytase (Aspergillus oryzae), FYT	Minimum	1 814 400	4 000 000
Endo-1,4-B-glucanase (T. longibrachiatum reesei), U	Minimum	120 960	266 667
Endo-1,3(4)-B-glucanase (T. longibrachiatum), U	Minimum	105 840	233 333

INGREDIENTS: Enzyme Preparation of Dried *Trichoderma longibrachiatum* Fermentation Extract, Rice Hulls, Enzyme Preparation of Dried *Aspergillus oryzae* Fermentation Extract, Calcium Carbonate.

DIRECTIONS FOR USE: Feed Victus™ Finisher at the rate of 0.75 pounds per ton (0.375 kilos per tonne) of complete poultry finisher feed.

One unit of phytase (FYT) is the amount of enzyme which liberates 1 mmol of inorganic phosphate per minute from sodium phytate at pH 5.5 and 37oC. One international unit (IU) of endo-1,4-beta-glucanase (cellulose) is the amount of enzyme which at pH 5.8 and 40oC, liberates 1 mmol glucose per min. One international unit (IU) of endo-1,3(4)-beta-glucanase (beta-glucanase) is the amount of enzyme which at pH 5.8 and 40oC, liberates 1 mmol glucose per min. One international unit (IU) of endo-1,4-beta-xylanase (xylanase) is the amount of enzyme which at pH 5.8 and 40oC, liberates 1 mmol xylose per min.

Danger

Hazard statements

May form combustible dust concentrations in air. H334 May cause allergy or asthma symptoms or breathing difficulties if inhaled. H350 May cause cancer.

Precautionary statements Prevention: P201 Obtain special instructions before use. P202 Do not handle until all safety precautions have been read and understood. P261 Avoid breathing dust/ fume/ gas/ mist/ vapours/ spray. P280 Wear protective gloves/ protective clothing/ eye protection/ face protection. P285 In case of inadequate ventilation wear respiratory protection.

Response: P304 + P341 IF INHALED: If breathing is difficult, remove victim to fresh air and keep at rest in a position comfortable for breathing. P308 + P313 IF exposed or concerned: Get medical advice/ attention.

Storage: P405 Store locked up.

Disposal: P501 Dispose of contents/ container to an approved waste disposal plant.

Contains: cellulase, glucanase, endo-1,3(4)-β-, Xylanase, endo-1,4-, phosphatase, acid, Quartz (SiO2)

For Use in Manufactured Feed Only

NP23108025

Net weight: 25 KG (55.115 POUNDS)

Manufactured for
DSM Nutritional Products
Parsippany, NJ

Date Manufactured:
00.00.0000

%
*F02475.100
*A0B44.44*SN_1.Y*FS
)%