



6.25" h

6.375" w

5.8125" w

5.8125" w

5.8125" w

5.8125" w

5.875" w

Virkon™ S Disinfectant and Virucide*

GREENHOUSES AND HORTICULTURE
Virkon™ S is effective against hard, non-porous environmental surfaces such as floors, walls, glasshouse structures, ventilation and other equipment, utensils, trays, and other containers, water systems, aqueous coolers, storage rooms, and vehicles in greenhouses and other horticultural settings prior to introduction or reintroduction of plants, seeds, or soil. Not approved in California for use on ventilation and other equipment and water systems. It is not intended to directly affect agricultural production and must not be applied to plants, seeds, or soil. If necessary, remove or cover these items prior to use of the product.

For swine and equine:
1. Sweep and remove all plant debris. Use power sprayer to wash all surfaces to remove loose dirt.
2. Use a dilution of 1:100 or 1.3 oz. Virkon™ S per gallon of clean water. Use a dilution of 1:50 or 2.6 oz. per gallon of clean water if surfaces that are to be treated have not been pre-cleaned with water to remove organic deposits. Not approved in California for use at 1:50 dilution on surfaces that have not been pre-cleaned with water to remove organic deposits.
3. Apply solution with mop, sponge, power sprayer or fogger to thoroughly wet all surfaces. Allow surfaces to remain visibly wet for a period of 10 minutes.
4. Heavy growth of algae or fungi may have to be scrubbed off following application.
5. Reapply as often as needed for control.

For clean non-porous hard surfaces
Pots, flats, trays: Use a dilution of 1:100 or 1.3 oz. per gallon of clean water. Soak tools to ensure complete coverage.
Walkways: Sweep and remove all plant debris. Use power sprayer to wash all surfaces to remove loose dirt. Use a dilution of 1:100 or 1.3 oz. of Virkon™ S per gallon of clean water. Use a dilution of 1:50 or 2.6 oz. of Virkon™ S per gallon of clean water if surfaces that are to be treated have not been pre-cleaned with water to remove organic deposits.

For evaporative coolers: Not an approved use in California. Treat existing algae and slime-contaminated surfaces with a 1:100 dilution of Virkon™ S. Treat cooler water every week with a dilution of 1:200 or 0.65 oz. of Virkon™ S for every gallon of cooler water.
Virkon™ S may also be used to descale irrigation tanks and lines. Not an approved use in California. Run a 1% solution through the system or soak equipment in a 1% solution. Let stand for ten minutes (allow surfaces to remain visibly wet for a period of 10 minutes) and flush system with clean water after treatment.

EMERGENCY/DISEASE CONTROL ANIMAL HEALTH
Disinfection helps to reduce cross-contamination of OIE List A organisms including Newcastle Disease Virus, Foot and Mouth Disease Virus, African Horse Sickness Virus, Vesicular Stomatitis Virus, Classical Swine Fever Virus, (Hog Cholera Virus), African Swine Fever Virus, Highly Pathogenic Avian Influenza Virus, Swine Vesicular Disease Virus, and Mycoplasma mycoides (Contagious Bovine Pleuropneumonia).
Disinfection helps to reduce cross-contamination of OIE List A organisms including Newcastle Disease Virus, Foot and Mouth Disease Virus, African Horse Sickness Virus, Vesicular Stomatitis Virus, Classical Swine Fever Virus, (Hog Cholera Virus), African Swine Fever Virus, Highly Pathogenic Avian Influenza Virus, Swine Vesicular Disease Virus, and Mycoplasma mycoides (Contagious Bovine Pleuropneumonia).
Not approved for this use in California.
Use a 1% solution of Virkon™ S to clean and disinfect hard non-porous surfaces associated with agricultural facilities and equipment, military facilities and equipment, sport facilities and equipment, port facilities and equipment, rail facilities and equipment, quarantine facilities and equipment, slaughter facilities and equipment, and other shipping facilities and equipment where animals or soils suspected of harboring Foot and Mouth Disease Virus might have been previously present.
Within these facilities, treated objects include but are not limited to vehicles, farm equipment (including tractors, plowing shares, cans and trucks, farm engines, harvesters, balers, mowers, tillers and slaughter machinery), military equipment (including tanks and troop carriers), and shipping equipment (pallets, bins and containers).
On surfaces that have been pre-cleaned with water and soap or detergent to remove organic deposits, spray Virkon™ S at 1% solution to disinfect and clean walls, ceilings, floors, decks, container surfaces,

Virkon™ S Disinfectant and Virucide*

USES IN FACILITIES USED FOR TEMPORARY CONFINEMENT OF ANIMALS
Use a 1% solution of Virkon™ S to clean and disinfect hard, non-porous surfaces associated with facilities used for the temporary confinement of animals. Sites may include, but are not limited to, barns, sheds, stables, pens, cages, and associated access alleys or walkways.
Virkon™ S may also be used to clean and disinfect equipment related to the maintenance of animals found at fairs, exhibitors, animal auction yards, animal show/boarding facilities, or other similar agricultural facilities designed for the temporary housing of animals.
To ensure that Virkon™ S does not come in direct contact with animals, feed, or water, remove animals from treatment site and either remove feed and water apparatus, or ensure precise application of Virkon™ S to hard, non-porous surfaces. Virkon™ S may only be applied to hand-drenched sprayers, sponges or other absorbent materials.
Do not allow Virkon™ S to pool on surfaces that may be within reach of animals. Do not allow Virkon™ S to come into direct contact with people. Allow Virkon™ S to completely dry prior to housing animals, using equipment, or allowing people to contact treated sites.

DISINFECTION LIMITED TO SPECIFIC AND KNOWN DISEASE ORGANISM PCV2 Virus*
Not approved for this use in California.
The instructions above call for use of a 1% solution for general disinfection. However, Virkon™ S is an effective disinfectant for hard, non-porous surfaces against the following organisms when the pathogen is limited to the organisms below. Virkon™ S may be used at the following dilution rate:

Organism	Dilution Rate	Oz./Gal.
PCV2 Virus (PMM5)	1:200	0.7

REDUCE CONTACT TIME (1 MINUTE) DISINFECTION LIMITED TO SPECIFIC AND KNOWN DISEASE CANINE PARVOVIRUS:
Not approved for this use in California.
The instructions above call for use of a 1% solution for general disinfection at a contact time of 10 minutes. However, Virkon™ S is an effective disinfectant for hard, non-porous surfaces against the following organisms with a 1 minute contact time when used at a dilution of 1:75. If the pathogen is limited to the organism below, Virkon™ S may be used at the following dilution rate:

Organism	Dilution Rate	Oz./Gal.
CANINE PARVOVIRUS	1:75	1.7

A 1.75 dilution is equal to 1.5 Scoops Virkon™ S in 1 Gallon of water.

Virkon™ S Disinfectant and Virucide*

REDUCED CONTACT TIME, COLD TEMPERATURE (5°C), AND/OR INCREASED DILUTION FOR DISINFECTION LIMITED TO SPECIFIC AND KNOWN DISEASES:
Not approved for this use in California.
The instructions above call for use of a 1% solution for general disinfection at a contact time of 10 minutes at room temperature. However, Virkon™ S is an effective disinfectant for hard, non-porous surfaces against the following organisms when the pathogen is limited to one of the organisms below. Virkon™ S may be used when the corresponding conditions are also applied:

Organism	Dilution Rate	Oz./Gallon	Contact Time (min)	Temperature (°C)
AAVAV ADENOVIRUS Type 2	1:200	0.7	10	5
AAVAV INFECTIOUS BRONCHITIS	1:200	0.7	1	5
AAVAV INFLUENZA virus	1:200	0.7	1	5
AAVAV INFLUENZA virus	1:600	0.215	10	RT
INFECTIOUS LARYNGOTRACHEITIS VIRUS	1:200	0.7	10	RT
MAREK'S DISEASE VIRUS	1:200	0.7	10	5
PDV	1:200	0.7	1	5
PDV	1:600	0.215	10	5
PRRS	1:600	0.215	10	5
PORCINE ROTAVIRUS	1:600	0.215	10	5

(RT = Room temperature = 20-22°C)
Instructions for making a 1:600[†] Dilution
Quantity of Water 1:600 Solution^{††}
10 Gallons 2.2 ounces / 2 Scoops
50 Gallons 11.0 ounces / 10 Scoops
1 measuring cup = 1 scoop = 1.3oz.
^{††} Not an approved use in California.

Virkon™ S Disinfectant and Virucide*

Using the Hydroblamer 481 (See 2) - How to Make a Concentrated Stock Solution
10lb/Pail of Virkon™ S = 1:600 oz.
(Not approved for this use in California)

Dilution Factor (DF)	Hydroblamer Tip Color	1:100 Dilution Rate (1.3 oz/gal)	1:600 Dilution Rate [†] (0.22oz/gal)
Calculation		$1.3 \text{ oz.} \times DF = \text{Volume of sprayer in gallons} \times \# \text{ of scoops} = \text{oz. Virkon™ S} / 1.3 \text{ oz.}$	$0.22 \text{ oz.} \times DF = \text{Volume of sprayer in gallons} \times \# \text{ of scoops} = \text{oz. Virkon™ S} / 1.3 \text{ oz.}$
1:20	No Tip	11.7 oz Virkon™ S 9 scoops of Virkon™ S	1.88 oz Virkon™ S 1.5 scoops of Virkon™ S
1:41	White	13.65 oz Virkon™ S 10.5 scoops of Virkon™ S	2.31 oz Virkon™ S 1.8 scoops of Virkon™ S
1:61	Blue	15.6 oz Virkon™ S 12 scoops of Virkon™ S	2.64 oz Virkon™ S 2 scoops of Virkon™ S
1:81	Tan	N/A	2.97 oz Virkon™ S 2.3 scoops Virkon™ S
3:21	Orange	N/A	5.28 oz Virkon™ S 4 scoops Virkon™ S
6:41	Purple	N/A	10.56 oz Virkon™ S 8.1 scoops Virkon™ S

[†]The 1:600 dilution is only approved for viruses of Avian Influenza Virus, Porcine Epidemic Diarrhea virus, Porcine Reproductive and Respiratory Syndrome, and Porcine Rotavirus. Please consult product label and use directions for 1:600 dilution for specified and known organisms.
Convert your Hydroblamer from ounces to gallons
32 oz = 0.25 gallon 96 oz = 0.75 gallon
64 oz = 0.5 gallon 128 oz = 1 gallon

Manufactured for: Lanxess Corporation, 111 Rittenhouse Park Drive, Pittsburgh, PA 15275
For information: US/Canada (800) LANXESS International +1 412 839 1000

©2011 LANXESS. Virkon™ S, LANXESS, the LANXESS Logo and any associated logos are trademarks or registered trademarks of LANXESS. Disinfectant and Virucide are trademarks of LANXESS. All trademarks are registered in many countries worldwide.

Net Contents: 22lb (9.98kg)
RSL_31/VHS/10b/USA/18.03.21/2

Virkon™ S Disinfectant and Virucide*

For use in cleaning and disinfecting industrial, animal and agricultural facilities
Powder Form
Effective against Viruses*, Bacteria, and Fungi
ACTIVE INGREDIENTS
Potassium peroxymonosulfate 21.41%
Sodium chloride 1.50%
OTHER INGREDIENTS
77.09%
Total 100.00%
Equivalent to 9.75% Available Chlorine
KEEP OUT OF REACH OF CHILDREN DANGER/PELIGRO
See Inside Booklet for Additional Precautions and explanatory notes
EPA REG. NO. 39967-137 EPA EST. NO. 57787-MI-003
FIRST AID IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for further treatment advice. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a Poison Control Center or doctor for further treatment advice. **IF SWALLOWED:** Call a Poison Control Center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. **HOT LINE NUMBER For 24-hour emergency information on this product, call 866-672-6358.** Have the product container or label with you when calling a poison control center or doctor or going for treatment. **NOTE TO PHYSICIAN:** Possible mucosal damage may contraindicate the use of gastric lavage.

Net Contents: 22lb (9.98kg)
RSL_31/VHS/10b/USA/18.03.21/2

Virkon™ S Disinfectant and Virucide*

PRECAUTIONARY STATEMENTS
DANGERS TO HUMANS AND DOMESTIC ANIMALS
HAZARD: Irritant is corrosive. Causes irreversible eye damage or skin burns. Harmful if swallowed or absorbed through the skin. Do not get in eyes, on skin or on clothing. Wear goggles (or face shield). Wear protective clothing (long sleeve shirt and long pants, socks plus shoes and chemical resistant gloves such as water proof gloves). Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash thoroughly before reuse.
Corrosive statement refers to powder only in use-diluted solution.
ENVIRONMENTAL HAZARDS
Do not discharge effluent containing the product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewerage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.
BROAD SPECTRUM DISINFECTANT:
Virkon™ S is effective against numerous microorganisms affecting animals: viruses*, gram positive and gram negative bacteria, fungi, and mycoplasma (sp., gallipneumoniae & hyppneumoniae). Efficacy of the 1% solution against bacteria and viruses* was determined in the presence of 400 ppm [200 ppm in California] AOAC hard water and 5% organic material in most cases. The exceptions are noted with qualifiers, e.g., "no hard water," "1% soil load," and "use 2% solution."

STORAGE AND DISPOSAL
Do not contaminate water, food, or feed by storage and disposal. Do not store this product with other chemicals.
PESTICIDE STORAGE: For pails, jugs, bottles: Store in a cool dry place in tightly closed container. Keep out of reach of children. Always replace the cap. Do not get in eyes, on skin or on clothing. Wear goggles (or face shield). Wear protective clothing (long sleeve shirt and long pants, socks plus shoes and chemical resistant gloves such as water proof gloves). Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash thoroughly before reuse.
PESTICIDE DISPOSAL: Pesticide wastes may be hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.
CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Triple-rinse container (or equivalent) promptly after emptying. Triple-rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Four rinses into application equipment and a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or follow the state and local authorities, by burning if burned, stay out of smoke.

Virkon™ S Disinfectant and Virucide*

EFFECTIVE AGAINST THE FOLLOWING ORGANISMS: ANIMAL AND ZOONOTIC PATHOGENS

BACTERIA	Moraxella bovis (Pink Eye)	Anthrax	Mandri-Virus Virus
<i>Bacillus cereus</i>	<i>Mycoplasma mycoides</i>	<i>Bovine Pasteurella Virus</i>	<i>Marx's Disease Virus</i>
<i>Bacillus anthracis</i>	<i>Pseudomonas mallei</i> (Glanders)	<i>Bovine Pseudocowpox Virus</i>	<i>Mouse Parvovirus</i>
<i>Clostridium perfringens</i>	<i>Staphylococcus hyicus</i>	<i>Bovine Viral Diarrhea Virus</i>	<i>PCV2 Virus (PMM5)</i>
<i>Clostridium botulinum</i>	<i>Streptococcus equi</i> (Strangles)	<i>Canine Distemper Virus</i>	<i>Porcine Epidemic Diarrhea Virus (PEDV)</i>
<i>Escherichia coli</i>	<i>Taylorella equigenitalis</i>	<i>Canine Parainfluenza Virus</i>	<i>Porcine Reproductive and Respiratory Syndrome Virus (PRRS)</i>
<i>Klebsiella pneumoniae</i>	<i>Trichinella spiralis</i>	<i>Canine Parvovirus</i>	<i>Porcine Rotavirus</i>
<i>Mycoplasma gallisepticum</i>	<i>Trichinella spiralis</i>	<i>Canine Adenovirus</i>	<i>Pseudorabies Virus (Aujeszky's Disease) (no hard water)</i>
<i>Mycoplasma hyopneumoniae</i>	<i>Trichinella spiralis</i>	<i>Canine Distemper Virus</i>	<i>Swine Vesicular Disease Virus</i>
<i>Pasteurella multocida</i>	<i>Trichinella spiralis</i>	<i>Canine Parvovirus</i>	<i>Swine Vesicular Disease Virus</i>
<i>Pseudomonas aeruginosa</i>	<i>Trichinella spiralis</i>	<i>Equine Herpes Virus (Type 1)</i>	<i>Transmissible Gastroenteritis Virus (TGE) (no hard water)</i>
<i>Salmonella enterica</i>	<i>Trichinella spiralis</i>	<i>Equine Herpes Virus (Type 2)</i>	<i>Turkey Herpes Virus</i>
<i>Salmonella typhimurium</i>	<i>Trichinella spiralis</i>	<i>Equine Influenza Virus (Type A)</i>	<i>Turkey Coronavirus</i>
<i>Staphylococcus aureus</i>	<i>Trichinella spiralis</i>	<i>Feline Calicivirus</i>	<i>Turkey Rhinotracheitis Virus</i>
<i>Staphylococcus epidermidis</i>	<i>Trichinella spiralis</i>	<i>Feline Parvovirus</i>	<i>Vesicular Stomatitis Virus</i>
<i>Streptococcus pyogenes</i>	<i>Trichinella spiralis</i>	<i>Feline Rhinotracheitis Virus</i>	
<i>Streptococcus</i>	<i>Trichinella spiralis</i>	<i>Feline Infectious Virus</i>	

Not approved in California for use against the following bacteria:
Bordetella avium
Adenovirus Pneumonia
Filatulus whartoni (Pill Evils)
Haemophilus somnus
Helicobacter pylori
Leptospira canicola
Listeria monocytogenes

Not approved in California for use against the following viruses:
African Horse Sickness Virus
Alphacoronavirus Virus (detected with 1% soil load and 342 ppm hard water)
Rhinotracheitis Virus (no hard water)

Virkon™ S Disinfectant and Virucide*

Virkon™ S DILUTION CHART
is a solution of Federal Use. Use this product in a manner inconsistent with its labeling.
is a solution of Federal Use. Use this product in a manner inconsistent with its labeling.
is a solution of Federal Use. Use this product in a manner inconsistent with its labeling.

Quantity of Water	0.5% Solution*	1% Solution	2% Solution
1 Quart	0.15 ounces	0.3 ounces	0.7 ounces
1 Gallon	0.65 ounces / 1/2 Scoop	1.3 ounces / 1 Scoop	2.7 ounces / 2 Scoops
10 Gallons	6.7 ounces / 5 Scoops	13.4 ounces / 10 Scoops	26.7 ounces / 20 Scoops
50 Gallons	33.4 ounces / 25 Scoops	66.8 ounces / 50 Scoops	133.5 ounces / 100 Scoops

^{††}The 0.5% solution currently is not approved for use in California.
Measuring cup provided.
1 measuring cup = 1 scoop = 1.3oz

DIRECTIONS FOR USE
is a solution of Federal Use. Use this product in a manner inconsistent with its labeling.
is a solution of Federal Use. Use this product in a manner inconsistent with its labeling.
is a solution of Federal Use. Use this product in a manner inconsistent with its labeling.

GENERAL INSTRUCTIONS - POULTRY AND FARM PREMISES
1. Remove all poultry or other animals and feed from pens, trucks or other vehicles, coops, crates or other enclosures.
2. Remove all litter droppings and manure from floors, walls and surfaces of barns, pens, stalls, crates and other facilities and fixtures occupied or surrounded by poultry or other animals.
3. Empty all troughs, racks, and other feeding and watering appliances.
4. Saturate surface with the use-diluted disinfecting solution. Allow surfaces to remain visibly wet for a period of 10 minutes.
5. Immerse all halteres, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and cups used for emptying litter and manure.
6. Wash buildings, cans, boots, coats, and other closed spaces. Do not house poultry or livestock or equipment until completely dry.
7. Turkeys should be treated necks, mangers, troughs, automatic feeders, fountains, and waters with soap or detergent, and rinse with potable water before reuse.

Virkon™ S Disinfectant and Virucide*

BOOT DISINFECTION
Virkon™ S at a 1% solution can be used for cleaning and disinfecting boots.
Boot Bath and Tray
Place boot bath or tray at entrance to reduce cross contamination on treated surfaces in poultry and farm premises. Use suitable containers and water containment areas.
1. Make a 1% solution of Virkon™ S by filling a suitable container with water and Virkon™ S according to the dilution table. Stir thoroughly until fully dissolved. Change the solution every 2 to 3 days or when it becomes visibly soiled.
2. Before application of Virkon™ S solution, rinse or brush footwear surfaces including soles to remove visible and heavy soil deposits.
3. Step into the container with boots on and use a clean brush to clean boots with disinfectant solution.
4. Wash boots thoroughly paying close attention to the soles. Allow surfaces to remain visibly wet for a period of 10 minutes.
5. Virkon™ S should only be used on water proof or non-porous footwear. Do not use on footwear that is damaged or permeable by water. Do not use on footwear that is damaged or permeable by water.

BROILER/BREEDER HOUSES: Follow General Instructions to remove poultry and pre-clean area to be treated. Spray floors and walls with Virkon™ S at 1% solution. Treated surfaces should remain visibly wet for duration of specified contact time. Thoroughly wash waterers and feeders with a 1% solution of Virkon™ S. Allow surfaces to remain visibly wet for a period of 10 minutes. After contact for 10 minutes, rinse with potable water. Do not house poultry or use equipment until treatment has dried.
PROCESSING PLANTS: Spray Virkon™ S at 1% solution to disinfect and clean walls, ceilings and floors. Saturate hard, non-porous surfaces with the use-diluted disinfection solution. Allow surfaces to remain visibly wet for a period of 10 minutes. Allow surfaces to air dry.

SWINE PRODUCTION
CONTROLS: Bacteria of *Actinobacillus pleuropneumoniae* and *Clostridium perfringens*, Foot and Mouth Disease Virus, Fungus of *Trichophyton interdigitale* (2%). **Not approved in California for use against the following organisms:** Viruses of Hog Cholera, Swine Vesicular Disease Virus, Porcine Epidemic Diarrhea Virus (PEDV), Porcine Reproductive and Respiratory Syndrome Virus (PRRS), Porcine Rotavirus, Pseudorabies Virus, Rotaviral Diarrhea Virus, Semeca Valley Virus (Slovakian Swine Fever Virus, Fungus of *Fusarium moniliforme*, and *Staphylococcus hyicus* (Greasy Pig) and *Trichophyton interdigitale* (2%).
General Instructions to remove swine and pre-clean area to be treated. Use a 1% solution of Virkon™ S for cleaning and disinfecting kennels, utility rooms, breeder houses, processing plants, and agricultural production equipment such as trucks, water/footwear (such as rubber boots), and associated livestock equipment and restrainments. Saturate surfaces with a 1% solution of Virkon™ S with a cloth, mop, mechanical spray or sponge. Allow surfaces to remain visibly wet for a period of 10 minutes. Allow surfaces to air dry. Rinse waterers and feeders with potable water before reuse.

Virkon™ S Disinfectant and Virucide*

EQUINE PRODUCTION
BROAD SPECTRUM EQUINE DISINFECTANT/DETERGENT/WASH FOR CLEANING AND DISINFECTING STABLES AND EQUIPMENT
CONTROLS: Viruses of Equine herpes Type 1 and Type 2 and Equine influenza. Bacteria of *Bordetella pertussis*, *Clostridium perfringens*, *Dermatophilus complexus* (2%), *Pseudomonas aeruginosa*, *Salmonella typhimurium* and *Staphylococcus aureus*. Fungus of *Trichophyton interdigitale* (at 2%).
Not approved in California for use against the following organisms: Fungus of *Fusarium moniliforme*, Viruses of African Horse Sickness, Equine Viral Arteritis (Pink Eye), Coat Ewartemia, Myxomatosis, Rhinopneumonitis, Equine Contagious Abortion, Equine Trichinella, Equine Infectious Anemia (Swamp Fever), Adenovirus Pneumonia, Equine Influenza (The Cough) and Rhinitis; Bacteria of *Clostridia*, *Diphtheria*, *Fibrotic Wither* (Poll Evil), *Taylorella equigenitalis*, *Yersinia bronchiseptica*, *Streptococcus equi* (Strangles) and *Pseudomonas mallei* (Glanders).
APPLICATIONS: For cleaning and disinfecting hard, non-porous surfaces, equipment, utensils and instruments in Veterinary practices, kennels, stables, trailers, etc.
USES: Stables, Horse Boxes, Box Stalls, Tack, Equipment, and Feed Rooms. Follow general instructions to remove animals from area to be treated. Thoroughly clean and dry (dry clean) surfaces, then wash the area manually or with pressure washer with a 1% Virkon™ S solution, saturating surfaces for a period of 10 minutes. Allow surfaces to remain visibly wet for a period of 10 minutes. Allow surfaces to air dry. Rinse waterers and feeders with potable water before reuse.
APPLICATIONS: Use a 1% solution of Virkon™ S as a "one step" cleaning and disinfecting procedure (Remove visible filth and soil deposits before application of the disinfecting/cleaning solution) for all surfaces, equipment, instruments, utensils and cages, caging systems, within or associated with Veterinary Medical Hospitals, animal infectious disease wards, quarantine areas, Humane Society facilities, laboratory animal quarters, grooming and boarding facilities, kennels, catteries and animal transportation vehicles.
Saturate surfaces with a 1% solution of Virkon™ S with a cloth, mop, mechanical spray or sponge. Allow surfaces to remain visibly wet for a period of 10 minutes. Allow surfaces to air dry. Rinse waterers and feeders with potable water before reuse. Do not remove metal objects in Virkon™ S for long periods.

5.875" w

5.8125" w

5.8125" w

5.8125" w

5.8125" w