

Wazine[®]

34

Turkey, Chicken and Swine Wormer

FDA Approved NADA 10-005



For use in drinking water for the removal of large roundworms (*Ascaridia spp.*) from turkeys, chickens and large roundworms (*Ascaris suum*) and nodular worms (*Oesophagostomum spp.*) from swine.

Active Drug Ingredient: Each 100 ml contains 34 grams piperazine base (present as sulfate).

Inert Ingredients: Water, color, flavoring, preservatives and stabilizer

DIRECTIONS:

For best results, remove water in the evening (except during hot weather). Then, early the next morning, give Wazine-34 in the drinking water. Provide medicated water only, distributed in waterers sufficient in number so that all birds or animals have access to water. The medicated water should be consumed in 1 day or less. Worming every 30 days helps break the large roundworm life cycle.

TURKEYS

Under 12 weeks of age: for each 1000 birds, use 10 fluid oz. (300 ml) of Wazine-34 in 20 U.S. gallons (76 liters) of drinking water.

Over 12 weeks of age: for each 1000 birds, use 20 fluid oz. (600 ml) of Wazine-34 in 40 U.S. gallons (150 liters) of drinking water.

Worm turkeys at 4 – 6 weeks of age and thereafter at 30 days or as needed.

CHICKENS

4 to 6 weeks of age: for each 1000 birds, use 5 fluid oz. (150 ml) of Wazine-34 in 10 U.S. gallons (38 liters) of drinking water.

Over 6 weeks of age: for each 1000 birds, use 10 fluid oz. (300 ml) of Wazine-34 in 20 U.S. gallons (76 liters) of drinking water.

Treat replacement pullets at 8, 12, and 16 weeks of age. Treat broilers at 4 weeks of age or as needed.

SWINE

For each 100 lb. (45.5 kg) of herd weight: give 1/2 fluid oz. (15 ml) of Wazine-34 per U.S. gallon (3.8 liters) of drinking water. Treat sows and gilts before breeding and up to 2 weeks before farrowing, treat boars at any time, and treat pigs 1 week after weaning and every 30 days thereafter.

CAUTION: Keep out of reach of children.
For animal use only.

NDC 15565-350-14
PL-2-341G-010906

Do not give to sick, feverish, underweight or physically weak birds or animals. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

WARNING: Do not medicate prior to slaughter as follows:

Turkeys – 14 days
Chickens – 14 days
Swine – 21 days

Do not use in chickens producing eggs for human consumption.
Restricted Drug – use only as directed

For use in automatic water proportioners: Mix 1 part Wazine-34 with 1 part water and set proportioner to deliver 1 fluid oz. (30 ml) per gallon of drinking water and give for 1 day.

One gallon (3.8 liters) of Wazine-34 provides a treatment for:

Turkeys: 12,800 up to 12 weeks of age, 6,400 over 12 weeks of age, or
Chickens: 25,600 4 to 6 weeks of age, 12,800 over 6 weeks of age, or
Swine: 1,024 25-lb. (11.4 kg) pigs, or 512 50-lb. (22.7 kg) pigs, or 256 100-lb. (45.5 kg) pigs, or 128 200-lb. (90.9 kg) pigs



Manufactured by
FLEMING LABORATORIES, INC.
Charlotte, NC 28234 USA

Net contents 3.785 L (1 U.S. Gallon)



Store above 32°F (0°C)